

# On the Determinants of Wroński in Linear Rings

by

M. KOWALSKI

*Presented by T. WAŻEWSKI on June 25, 1956*

**1.** In operational calculus [1] we consider the differential equations

$$(1) \quad x^{(n)} + a_{n-1}x^{(n-1)} + \dots + a_0x = 0,$$

in which  $a_j$  ( $j=0,1,\dots,n-1$ ) are given elements of a field of operators, and  $x$  is the unknown function assigning operators to real numbers. Equations (1) contain, as a particular case, ordinary differential equations — linear, homogeneous, with constant coefficients. Thus the results presented in this paper are valid also in the theory of ordinary differential equations.

**2.** Let  $C$  denote a commutative ring with unity and  $K$  any commutative linear ring on  $C$ . Assume that a linear endomorphism on elements of  $K$  was defined, which transforms  $x$  into  $x^{(1)}$  and in general  $x^{(n)}$  into  $x^{(n+1)}$  ( $n=1,2,\dots$ ). Let  $a_j \in C$  ( $j=0,1,\dots,n-1$ ),  $b_i \in C$  ( $i=0,1,\dots,k-1$ ). Suppose that we know the linearly independent solutions of the equation

$$L_fx \equiv x^{(n)} + a_{n-1}x^{(n-1)} + \dots + a_0x = 0,$$

which corresponds to the polynomial  $f(w) = w^n + a_{n-1}w^{n-1} + \dots + a_0$  and the linearly independent solutions of the equation

$$L_gx \equiv x^{(k)} + b_{k-1}x^{(k-1)} + \dots + b_0x = 0,$$

which corresponds to the polynomial  $g(w) = w^k + b_{k-1}w^{k-1} + \dots + b_0$ .

Let us form a third equation,  $L_hx \equiv L_f(L_gx) = 0$ . What are the solutions of equation  $L_hx = 0$ , and are they linearly independent? The answer to the second question is given by Theorem 2, valid in any linear space. The theorems on the determinants of Wroński have, incidentally, been proved and can also serve for the proof of linear independence of the

solutions of the equation  $L_h x = 0$ . The Theorems 1, 3 and 4 are valid in the ring  $K$ .

**3.** By the determinant of Wronski of the elements  $y_1, \dots, y_n$  we understand the determinant

$$\begin{vmatrix} y_1, \dots, y_1^{(n-1)} \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_n, \dots, y_n^{(n-1)} \end{vmatrix}.$$

**THEOREM 1.** If  $y_1, \dots, y_n$  are solutions of equation  $L_f x = 0$  of "order"  $n$ , and  $z_1, \dots, z_k$  are solutions of equation  $L_g x = 0$  of "order"  $k$ , both equations having unity as the "highest coefficient", then the determinant of Wronski  $W_h$  of the  $(n+k)$  elements  $y_1, \dots, y_n, z_1, \dots, z_k$  is equal to the product of the determinants of Wronski  $W_f$  of the elements  $y_1, \dots, y_n$  and  $W_g$  of elements  $z_1, \dots, z_k$  and the resultant  $R_{g,f}$  of polynomials  $g$  and  $f$ :

$$W_h = W_f \cdot W_g \cdot R_{g,f}.$$

**4.** Making use of the fact that  $ax = 0$  ( $a \in C$ ,  $x \in K$ ) only if  $a = 0$  or  $x = 0$ , it is easy to prove the following theorem:

**THEOREM 2.** If equations  $L_f x = 0$  and  $L_g x = 0$  with coefficients from an arbitrary \*) commutative ring  $C$  have a common solution  $x_1 \neq 0$ , then  $R_{f,g} = 0$  ( $R_{f,g}$  is the resultant of polynomials  $f$  and  $g$ ).

If  $C$  is a ring which has no divisors of zero \*\*), the vanishing of the resultant  $R_{f,g}$  is equivalent to the polynomials  $f, g$  having a common factor of positive degree. In particular, the discriminant of an irreducible polynomial in a ring which has no divisors of zero or in the field  $C$  is different from zero.

Denote by  $y_1, \dots, y_n$  solutions of the equation  $L_f x = 0$  and by  $z_1, \dots, z_k$  solutions of the equation  $L_g x = 0$ . If  $y_1, \dots, y_n, z_1, \dots, z_k$  are linearly dependent, but  $y_1, \dots, y_n$  are linearly independent and  $z_1, \dots, z_k$  are also independent, then the equations  $L_f x = 0$  and  $L_g x = 0$  have a common solution  $x_1 \neq 0$  and the resultant  $R_{f,g}$  vanishes.

**5.** It is known [2] that if  $x_i$  ( $i = 1, \dots, n$ ) satisfy the equation  $L_f x = 0$  corresponding to the polynomial  $f$ , and if the linear operation  $A$  satisfies the condition  $(Ax)^{(1)} = Ax^{(1)} + x$ , then the elements  $A^x x_i$  ( $x = 0, 1, \dots, k-1$ ;  $i = 1, \dots, n$ ) satisfy the equation  $L_{f^k} x = 0$ , corresponding to the  $k$ -th power of the polynomial  $f$ .

\*) The ring  $C$  does not necessarily contain a unity. Theorem 2 is valid for arbitrary polynomials  $f(w) = a_n w^n + \dots + a_0$ ,  $g(w) = b_k w^k + \dots + b_0$ .

\*\*) In the ring of continuous real functions with ordinary addition and multiplication let us take  $a \neq 0$ ,  $b \neq 0$  such that  $ab = 0$ ,  $a > 0$ ,  $b > 0$ . The polynomials  $f(w) = w^2 + a$ ,  $g(w) = w^2 + bw + a$  are irreducible but  $R_{f,g} = 0$ .



The following theorem expresses the determinant

$$W_f^k = \begin{vmatrix} x_1, & x_1^{(1)}, & \dots, & x_1^{(nk-1)} \\ \dots & \dots & \dots & \dots \\ x_n, & x_n^{(1)}, & \dots, & x_n^{(nk-1)} \\ Ax_1, & (Ax_1)^{(1)}, & \dots, & (Ax_1)^{(nk-1)} \\ \dots & \dots & \dots & \dots \\ Ax_n, & (Ax_n)^{(1)}, & \dots, & (Ax_n)^{(nk-1)} \\ A^{k-1}x_1, & (A^{k-1}x_1)^{(1)}, & \dots, & (A^{k-1}x_1)^{(nk-1)} \\ \dots & \dots & \dots & \dots \\ A^{k-1}x_n, & (A^{k-1}x_n)^{(1)}, & \dots, & (A^{k-1}x_n)^{(nk-1)} \end{vmatrix}$$

by the  $k$ -th power of the determinant

$$W_f = \begin{vmatrix} x_1, \dots, x_1^{(n-1)} \\ \dots & \dots & \dots \\ x_n, \dots, x_n^{(n-1)} \end{vmatrix}.$$

THEOREM 3. If  $x_1, \dots, x_n$  are solutions of the equation

$$L_f x \equiv x^{(n)} + a_{n-1}x^{(n-1)} + \dots + a_0x = 0,$$

and for those  $x_i$  ( $i=1, \dots, n$ ) the conditions

$$(2) \quad \begin{aligned} (Ax)^{(1)} &= Ax^{(1)} + x \\ x^{(\mu)} \cdot A^p x^{(\nu)} &= x^{(\nu)} \cdot A^p x^{(\mu)} \\ (p=1, \dots, k-1; \quad \mu, \nu=0, 1, \dots, nk-1) \end{aligned}$$

are satisfied, then

$$W_f^k = [1!2!\dots(k-1)!]^n \cdot R_{f,f'}^{k(k-1)/2} \cdot W_f^k,$$

where  $R_{f,f'}$  is the discriminant of the polynomial  $f$ .

The second of conditions (2) can, of course, be replaced by a somewhat stronger but simpler condition:  $A(xy) = (Ax)y$ .

**6.** An arbitrary polynomial  $F(w)$  with coefficients from a ring having no divisors of zero may be decomposed into factors irreducible in this ring.

$$F(w) = \prod_{j=1}^p f_j^{a_j}(w).$$

Let us take only those factors  $f_j(w)$  to which correspond equations having solutions  $x_{j1}, \dots, x_{jm_j}$  ( $j=1, \dots, q$ ), different from zero. By Theorem 2, from the linear independence of elements  $x_{j1}, \dots, x_{jm_j}$  we can deduce the linear independence of elements  $A^r x_{js}$  ( $r=0, 1, \dots, a_j-1$ ;  $j=1, \dots, q$ ;  $s=1, \dots, m_j$ ).

7. Suppose that  $F(w)$  can be decomposed into the factors

$$f_j(w) = w^{n_j} + a_{n_j-1}w^{n_j-1} + \dots + a_0$$

$$F(w) = c \prod_{j=1}^q f_j^{a_j}(w).$$

Applying successively Theorems 1 and 3, we obtain a theorem from which it may be deduced that the linear independence of the solutions  $x_{j1}, \dots, x_{jn_j}$  of the equations  $L_{f_j}x = 0$  implies the linear independence of  $A^r x_{js}$  ( $r = 0, 1, \dots, a_j - 1$ ;  $j = 1, \dots, q$ ;  $s = 1, \dots, n_j$ ).

THEOREM 4. *If the assumptions of Theorems 1 and 3 are satisfied, then*

$$W_F = \prod_{j=1}^q [1! 2! \dots (a_j - 1)!]^{n_j} \cdot \prod_{1 \leq \mu < \mu' \leq q} R_{f_\mu f_{\mu'}}^{\alpha_\mu \alpha_{\mu'}} \cdot \prod_{j=1}^q R_{f_j f_j'}^{a_j(a_j-1/2)} \prod_{j=1}^q W_{f_j}^{\alpha_j}.$$

8. Theorem 4 contains as a particular case the result of J. Mikusiński [3]. In order to obtain this result, it must be assumed that the polynomial  $F(w)$  can be decomposed into linear factors and that there exist elements  $x_j$ , different from zero, which satisfy equations  $x^{(1)} = w_j x$  ( $j = 1, \dots, q$ ),  $w_j$  being the roots of polynomial  $F(w)$  and the product of those  $x_j$  ( $j = 1, \dots, q$ ) being different from zero.

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES

#### REFERENCES

- [1] J. Mikusiński, *Sur les fondements du calcul opératoire*, Studia Mathematica **12** (1950), 41.
- [2] — *Sur les équations différentielles du calcul opératoire*, Studia Mathematica **12** (1951), 229.
- [3] — *Sur un déterminant*, Annales Soc. Pol. Math., **25** (1952), 27.

## Note of the Representation of Large Integers as Sums of Primes

by

YIN WEN-LIN

*Presented by W. SIERPIŃSKI on October 24, 1956*

Let  $K(n)$  denote the number of primes such that the integer  $n$  is expressible as a sum of  $K(n)$  but not as a sum of less than  $K(n)$  primes. Shapiro and Warga [1] proved that  $K(n) \leq 20$  for  $n > N$ . In this note it is shown that a slight improvement on their result is obtainable; actually we can prove that  $K(2n) \leq 18$  and  $K(2n+1) \leq 17$  for  $n > N$ . A sketch of the proofs is given below.

Let  $\Sigma$  be the set of all non-negative even integers and  $A \subset \Sigma$ . We define the "even" asymptotic density of  $A$  by

$$\delta_{\Sigma}^*(A) = \lim_{n \rightarrow \infty} \frac{A(2n)}{n},$$

where  $A(x)$  denotes, as usual, the number of integers in  $A$  not exceeding  $x$ . Analogous to a known property of the asymptotic density [2] is the relation

$$(1) \quad \delta_{\Sigma}^*(A+B) \geq \delta_{\Sigma}^*(A) + \frac{k-1}{k} \delta_{\Sigma}^*(B),$$

provided that  $A, B \subset \Sigma$ ,  $\delta_{\Sigma}^*(A) + \delta_{\Sigma}^*(B) \leq 1$  and  $B$  contains a chain of  $k$  consecutive even integers.

Further, we may define the "even" average order in a way analogous to the ordinary average order. Then, if  $0 \in B \subset \Sigma$ ,  $A \subset \Sigma$ ,  $\delta_{\Sigma}^*(A) = a$  and  $\lambda^*$  is the even average order of  $B$ , we have

$$(2) \quad \delta_{\Sigma}^*(A+B) \geq a \left( 1 + \frac{1-a}{\lambda^*} \right),$$



which is also analogous to a known result [3]. A further analogue to the known result is the relation

$$(3) \quad A+B \sim \Sigma,$$

provided that  $A, B \subset \Sigma$  and  $\delta_{\Sigma}^*(A) + \delta_{\Sigma}^*(B) > 1$ , where “ $\sim$ ” denotes that the set on the two sides of (3) differ by only a finite number of elements.

The results (1), (2) and (3) can be justified by an argument of “isomorphism”.

We use  $p$  and  $p'$  to denote odd primes only. It is known by [1], that for  $M = \{p + p'\}$ ,

$$\lim_{x \rightarrow \infty} \frac{M(x)}{x} \geq \frac{1}{16}.$$

Now consider  $P_0 = \{p - 3\}$ , then  $\delta_{\Sigma}^*(P_0 + P_0) = \delta_{\Sigma}^*(M)$ . It is easily verified that  $2n \in P_0 + P_0$  for  $1 \leq n \leq 125$ . Hence by (4) and (1)

$$\delta_{\Sigma}^*(hM) = \delta_{\Sigma}^*(h(P_0 + P_0)) \geq \text{Min} \left\{ \frac{1}{8} + \left( 1 - \frac{1}{125} \right) \frac{h-1}{8}, 1 \right\},$$

where, of course,  $hA$  is an abbreviation for  $A + \dots + A$  ( $h$  terms). Thus,

$$\delta_{\Sigma}^*(8M) + \delta_{\Sigma}^*(M) = \delta_{\Sigma}^*(8(P_0 + P_0)) + \delta_{\Sigma}^*(2P_0) > 1.$$

From (3) we find that

$$\Sigma \sim 9M \sim 9(P_0 + P_0).$$

It follows that every sufficiently large even integer can be expressed as a sum of at most 18 primes and also that the even average order of  $P$  is  $\lambda^* \leq 18$ . So, in view of (2), we have

$$\delta_{\Sigma}^*(M + P_0) \geq \frac{1}{8} \left( 1 + \frac{1 - \frac{1}{8}}{18} \right),$$

and hence

$$\delta_{\Sigma}^*(7M) + \delta_{\Sigma}^*(M + P_0) > 1 + \frac{7}{8 \cdot 8 \cdot 18} - \frac{6}{8 \cdot 125} > 1^*.$$

Therefore, by (3),

$$\Sigma \sim 8M + P_0$$

A moment of consideration shows that every large odd integer is expressible as a sum of at most 17 primes.

It should be noted that our result is far weaker than what can be deduced from the famous Goldbach-Vinogradov theorem, namely  $K(n) \leq 4$  for sufficiently large  $n$ . Our proof, however, is elementary in the sense

---


\*) The same result can also be obtained by using an analogue of Erdős' result instead of that of Kasch's.

that we make no use of any deep analytic method or theorem. Besides what can be seen from the note itself, our proof depends indirectly upon Selberg's sieve method which may also be regarded as elementary.

PEKING UNIVERSITY

# REFERENCES

- [1] H. N. Shapiro and J. Warga, *On representation of large integers as sums of primes I.*, Comm. pure Appl. Math. **3** (1950), 153.
- [2] H. H. Ostmann, *Vereinerte Lösung der asymptotischen Dichtaufgabe*, J. Reine Angew. Math. **187** (1950), 183.
- [3] F. Kasch, *Abschätzung der Dichte von Summenmengen*, Math. Zeit. **64** (1956), 243.
- [4] H. Rohrbach, *Einige neuere Untersuchungen über die Dichte in der additiven Zahlentheorie*, Jber. Deutschen Math. Verein **43** (1938), 199.
- [5] P. Erdős, *On the asymptotic density of the sum of two sequences one of which forms a basis for the integers II.*, Trav. Inst. Math. Tbilissi **3** (1938), 217.



Digitized by the Internet Archive  
in 2025



# On Algebraic Structures and Homotopy Invariants

by

S. C. CHANG

*Presented by K. BORSUK on October 9, 1956*

Let  $X$  be a geometric object. According to the geometric properties of  $X$  a lot of algebraic structures are constructed to "realise" them. For instance, in a Euclidean space we have parallel lines in a given direction, — this property evidently determines a translation group (in the given direction) whose trajectories are the given parallel lines. It is interesting in history to study the algebraic properties of the algebraic structures associated with a geometric object. This is the fundamental ideal of algebraic topology.

In 1935, W. Hurewicz [1] introduced homotopy types of topological spaces. Afterwards J. H. C. Whitehead defined cohomology rings [2] and cohomology systems [3]. He produced a 1-1 correspondence between isomorphism classes of cohomology rings and the homotopy types of simply connected 4-dimensional polyhedra and then he simplified his argument to establish a similar correspondence between isomorphism classes of cohomology systems and  $A_n^2$ -polyhedra ( $n \geq 3$ ). According to Whitehead an arcwise connected polyhedron,  $K$ , is called an  $A_n^2$ -polyhedron if  $\dim K \leq n+2$  and  $\pi_r(K) = 0$ ,  $r = 1, \dots, n-1$ . We assume  $n \geq 3$ . Let  $H^n, H^{n+1}, H^{n+2}$  be additive groups, each of which has a finite number of generators,  $H^n$  being free. A new group,  $H^n(2)$ , is defined as the direct sum \*)

$$H_2^n + \Delta^*({}_2H^{n+1}),$$

where  $\Delta^*({}_2H^{n+1})$  is the image of  ${}_2H^{n+1}$  under an isomorphism  $\Delta^*$ . Here  $\Delta^*$  is determined modulo an arbitrary homomorphism

$${}_2H^{n+1} \rightarrow H_2^n.$$

---

\*) Let  $G$  be an additive Abelian group then  $G_2 = G - 2G$ , and  ${}_2G$  is the subgroup of  $G$  which consists of all the elements  $g$ , such that  $2g = 0$ .

In other words, the isomorphism  $\Delta^*$  is not determined uniquely, but there is a uniquely determined homomorphism

$$\Delta: H^n(2) \rightarrow {}_2H^{n+1},$$

such that  $\Delta\Delta^*=1$  and  $\Delta^{-1}(0)=H_2^n$ . Furthermore, a homomorphism  $\mu: H^r \rightarrow H^r(2)$ , ( $r=n, n+2$ ), onto  $H_2^r$  is defined in the usual way. Finally, there is given a homomorphism  $\gamma^*: H^n(2) \rightarrow H^{n+2}(2)$ , which may be arbitrary. Then  $H^n, H^{n+1}, H^{n+2}, H^n(2), H^{n+2}(2)$ , together with  $\Delta, \mu$  and  $\gamma^*$  constitute an  $A_n^2$ -cohomology system. This is the algebraic cohomology system of J. H. C. Whitehead. On the other hand, if an  $A_n^2$ -polyhedron,  $K$ , is given, we have the integral cohomology groups  $H^r$  ( $r=n, n+1, n+2$ ) and the cohomology groups  $H^r(2)$ , ( $r=n, n+2$ ) with integers reduced mod. 2 as coefficients, and we take the natural homomorphism  $\mu, \Delta = \frac{1}{2}\delta$  and finally  $\gamma^*$  as the Steenrod square [4]  $\gamma^*x = x \cup_{n-2} x_e H^{n+2}(2)$ , where  $x_e H^n(2)$ . Therefore a cohomology system is associated with  $K$ . A homomorphism  $f$  between two cohomology systems contains five homomorphisms of the corresponding groups with a restriction that  $f$  commutes with  $\mu, \Delta$  and  $\gamma^*$ . Particularly  $f$  is called a  $(\mu, \Delta)$ -homomorphism if  $f$  commutes with  $\mu$  and  $\Delta$  only.

Now cohomology groups are invariants of homotopy type. How many different homotopy types of  $A_n^2$ -polyhedra may there be if their cohomology groups are assigned? In other words, we are required to study the algebraic properties of cohomology systems so as to find the necessary and sufficient conditions that two cohomology systems are isomorphic if their groups are isomorphic. We discovered [5] new numerical invariants called secondary torsions and reached the following result:

**THEOREM.** *The homotopy type of an  $A_n^2$ -polyhedron ( $n > 2$ ) is determined by its Betti numbers, torsions and secondary torsions.*

In this theorem Betti numbers, torsions and secondary torsions become a *complete* and *independent* system of invariants of the homotopy type of  $A_n^2$ -polyhedra.

By an elementary  $A_n^2$ -polyhedron we mean one of the following kinds:

- (1)  $B_1^r = S^r$  ( $r=n, n+1, n+2$ );
- (2)  $B_2(\sigma) = S^n \cup e^{n+1}$ , where  $e^{n+1}$  is attached to  $S^n$  by a map  $f: \dot{E}^{n+1} \rightarrow S^n$  of degree  $\sigma$ , a power of a prime;
- (3)  $B_3(\tau) = S^{n+1} \cup e^{n+2}$ , where  $e^{n+2}$  is attached to  $S^{n+1}$  by a map  $f: \dot{E}^{n+2} \rightarrow S^{n+1}$  of degree  $\tau$ , a power of a prime;
- (4)  $B_4 = S^n \cup e^{n+2}$ , where  $e^{n+2}$  is attached to  $S^n$  by an essential map  $f: \dot{E}^{n+2} \rightarrow S^n$ ;
- (5)  $B_5(2^p) = S^n \cup e^{n+1} \cup e^{n+2}$ , where  $e^{n+1}$  is attached to  $S^n$  by an essential map  $f: \dot{E}^{n+2} \rightarrow S^n$ ;



- (6)  $B_6(2^a) = S^n \cup S^{n+1} \cup e^{n+2}$ , where  $S^n \cap S^{n+1} = x_0$ , a point, and  $e^{n+2}$  is attached to  $S^n \cup S^{n+1}$  by a map  $f: \dot{E}^{n+2} \rightarrow S^n \cup S^{n+1}$  of the form  $a + b$ , where  $a$  denotes an essential map of  $\dot{E}^{n+2}$  onto  $S^n$  and  $b$  maps  $\dot{E}^{n+2}$  onto  $S^{n+1}$  with degree  $2^a$ ;
- (7)  $B_7(2^p, 2^a) = B_6(2^a) \cup e^{n+1}$ . Attach  $e^{n+1}$  to  $S^n$  of  $B_6(2^a)$  by a map  $f: \dot{E}^{n+1} \rightarrow S^n$  with degree  $2^p$ .

Take a certain number of elementary polyhedra, choose a point on the sphere of lowest dimension of each elementary polyhedron and identify all these points to a new point  $p_0$ , thus forming an  $A_n^2$ -polyhedron,  $K$ , which is a normal form for the homotopy type of a given  $A_n^2$ -polyhedron.

By means of normal  $A_n^2$ -polyhedron  $\pi_{n+1}(K)$  is computed ([5], [6]), as a generalisation of the well-known Hurewicz theorem. Furthermore, P. J. Hilton [6] computed the  $(n+2)$ -dimensional homotopy group of  $A_n^2$ -polyhedron (if  $n \geq 3$ ) while the Borsuk-Spanier cohomotopy group  $\pi^n(K^{n+1})$ , ( $n > 3$ ),  $K^{n+1}$  being a given  $(n+1)$ -dimensional polyhedron, is expressed [5] by means of the Betti numbers, torsions and secondary torsions of a special  $A_n^2$ -polyhedron obtained from  $K^{n+1}$  by identifying its  $(n-1)$ -skeleton to a point.

In the definition of the cohomology ring the Pontrjagin square is involved in place of the Steenrod square. The algebraic classification of cohomology rings seems to be very difficult.

In 1950, J. H. C. Whitehead and myself [7] extended the secondary torsions associated with the  $A_n^2$ -polyhedron ( $n \geq 3$ ) to block invariants associated with a finite polyhedron. We [7] considered  $(\mu, \Delta, \gamma)$ -systems which consist of the whole family of groups  $H^n(m)$  ( $m=0, 2$ ) related by the homomorphisms  $\mu, \Delta, \gamma$ , where  $\gamma = \gamma_k^n: H^n(2) \rightarrow H^{n+k}(2)$  is supposed to be defined for every even  $k > 0$  and every  $n \geq k$ . To a given topological space a  $(\mu, \Delta, \gamma)$ -system is evidently determined by the geometrical method. Since a homotopy equivalence  $f: X \rightarrow Y$  induces an isomorphism of  $(\mu, \Delta, \gamma)$ -systems related to  $Y$  and  $X$ , algebraic invariants of the isomorphism class of  $(\mu, \Delta, \gamma)$ -systems must be invariants of the homotopy type. From this view-point many numerical homotopy invariants may be found if we are able to classify  $(\mu, \Delta, \gamma)$ -systems according to their isomorphism classes.

No doubt we may have an algebraic structure if we have a cohomology operation (or a set of cohomology operations) which commutes with the homomorphism between cohomology groups induced by a continuous mapping. Particularly, we take two Steenrod homomorphisms,  $\gamma_j^m: H^m(2) \rightarrow H^{m+j}(2)$ ,  $\gamma_k^n: H^n(2) \rightarrow H^{n+k}(2)$ , and construct a special  $(\mu, \Delta, \gamma)$ -system containing  $\gamma_j^m$ ,  $\gamma_k^n$  and their related cohomology groups  $H^r$  ( $r=m, m+1, m+j, m+j+1, n, n+1, n+k, n+k+1$ ) and  $H^s(2)$  ( $s=m, m+j, n, n+k$ ). In [7] it has been actually remarked that block invariants together with Betti numbers and torsions are not enough to characterise isomorphism classes of these special  $(\mu, \Delta, \gamma)$ -systems, if among the sets of groups



$(H^m, H^{m+1}, H^{m+j}, H^{m+j+1})$  and  $(H^n, H^{n+1}, H^{n+k}, H^{n+k+1})$  there are common groups. Now there are two extremities: In the simplest case there is one common group between the sets  $(H^m, H^{m+1}, H^{m+j}, H^{m+j+1})$  and  $(H^n, H^{n+1}, H^{n+k}, H^{n+k+1})$ , but in the most complicated case there exist three common groups. The isomorphism class of  $(\mu, \Delta, \gamma)$ -systems of the first case has been studied in [8]; block invariants, relative block invariants, Betti numbers and torsions are complete and independent sets of invariants. In the latter case we have  $j=k=2$  and  $n=m+1$ , which system plays an important role in the homotopy type of  $A_n^3$  ( $n \geq 3$ ) polyhedron and is called the  $A_n^3$ -system. In a recent paper [9] we discover new numerical invariants called characteristic coefficients and characteristic polynomials and have proved that isomorphism classes of  $A_n^3$ -systems are characterised by Betti numbers, torsions, block invariants, relative block invariants, characteristic coefficients and characteristic polynomials. All of these invariants are invariants of a given finite polyhedron.

In the end I remark that general steps to characterise  $(\mu, \Delta, \gamma)$ -systems are indicated in [8]. We expect to find complete and independent systems of invariants of homotopy types of polyhedra and determine normal forms of these types. Normal forms of polyhedra appear to be useful in several occasions.

PEKING UNIVERSITY

#### REFERENCES

- [1] W. Hurewicz, Proc. Akad. Sci. Amst. **38** (1935), 112, 521; *ibid.* **39** (1936), 117, 215.
- [2] J. H. C. Whitehead, Comm. Math. Helv. **22** (1949), 48.
- [3] — Ann. Soc. Math. Polon. **21** (1948), 176.
- [4] N. E. Steenrod, Ann. Math. **48** (1947), 290.
- [5] S. C. Chang, Proc. Royal Soc. (British), **202** (1950), 253.
- [6] P. J. Hilton, Proc. L. M. S. **1** (1951), 462.
- [7] S. C. Chang and J. H. C. Whitehead, Quart. J. Math. (Oxford), **2** (1951), 167.
- [8] S. C. Chang, Bull. Acad. Polon. Sci. Cl. III, **4** (1956), 113.
- [9] — unpublished.

## Sur la division des corps matériels en parties

par

H. STEINHAUS

Présenté le 19 Octobre 1956

Un corps  $Q$  est, par définition, une répartition de matière dans l'espace, donnée par une fonction  $f(P)$ ; on appelle cette fonction la *densité* du corps en question; elle est définie pour tous les points  $P$  de l'espace; elle est non-négative et mesurable. On suppose que l'ensemble caractéristique du corps  $E = \bigcup_P \{f(P) > 0\}$  est borné et de mesure positive; on suppose aussi que l'intégrale de  $f(P)$  sur  $E$  est finie: c'est la *masse* du corps  $Q$ . On considère comme identiques deux corps dont les densités sont égales à un ensemble de mesure nulle près.

En décomposant l'ensemble caractéristique d'un corps  $Q$  en  $n$  sous-ensembles  $E_i$  ( $i=1,2,\dots,n$ ) de mesures positives, on obtient une *division* du corps en question en  $n$  *corps partiels*; leurs ensembles caractéristiques respectifs sont les  $E_i$  et leurs densités sont définies par les valeurs que prend la densité du corps  $Q$  dans ces *ensembles partiels*. En désignant les corps partiels par  $Q_i$ , on écrira  $Q=Q_1+Q_2+\dots+Q_n$ . Quand on donne d'abord  $n$  corps  $Q_i$ , dont les ensembles caractéristiques sont disjoints deux à deux à la mesure nulle près, il existe évidemment un corps  $Q$  ayant ces  $Q_i$  comme autant de parties; on écrira  $Q_1+Q_2+\dots+Q_n=Q$ . Ces remarques suffisent pour expliquer la division et la composition des corps.

Le PROBLÈME de cette Note est la division d'un corps en  $n$  parties  $K_i$  ( $i=1,2,\dots,n$ ) et le choix de  $n$  points  $A_i$  de manière à rendre aussi petite que possible la somme

$$(1) \quad S(K,A) = \sum_{i=1}^n I(K_i, A_i) \quad (K \equiv \{K_i\}, \quad A \equiv \{A_i\}),$$

où  $I(Q,P)$  désigne, en général, le *moment d'inertie* d'un corps quelconque  $Q$  par rapport à un point quelconque  $P$ . Pour traiter ce problème élémentaire nous aurons recours aux lemmes suivants:

a)  $Q$  étant donné,  $I(Q, P)$  atteint un minimum parfait quand  $P$  devient identique avec le centroïde de  $Q$ .

b) Si  $(K, A)$  est la solution du problème, tout  $A_i$  est le centroïde de  $K_i$  ( $i=1, 2, \dots, n$ ).

c) Soient  $K_1$  et  $K_2$  deux corps partiels de  $Q$ , obtenus en coupant l'ensemble caractéristique  $E$  de  $Q$  en deux par un plan; soient  $A_1$  et  $A_2$  les centroïdes respectifs de  $K_1$  et  $K_2$ ; on aura

$$S(K, A) < I(Q, P)$$

quel que soit le point  $P$ .

d) Si  $(K, A)$  est la solution du problème, on a  $A_i \neq A_j$  pour  $i \neq j$  ( $i, j=1, 2, \dots, n$ ).

e) Soit  $\bar{D}$  le plus petit domaine convexe et fermé contenant l'ensemble caractéristique  $E$  de  $Q$ , et soit  $P'$  le point de  $\bar{D}$  le plus proche de  $P$ ; on aura  $I(Q, P) \geq I(Q, P')$ .

Le lemme a) résoud le problème pour  $n=1$ ; il est connu universellement; nous y parlons de minimum *parfait*, car il n'y a pas d'autres points donnant des valeurs extrêmes à  $I$ . Le lemme b) résulte de a). Le lemme c) résulte de ce que  $I(Q, P) = I(K_1, P) + I(K_2, P)$ , de a) et du fait que le centroïde d'un corps  $Q$  quelconque est toujours intérieur au domaine  $D = \bar{D} - F$  (on a défini  $\bar{D}$  dans e) et  $F$  est la frontière de  $\bar{D}$ ); on applique ce fait aux corps  $A_1$  et  $A_2$ . Le lemme d) résulte de l'application de c) au corps  $K_i + K_j$ . Le lemme e) est fondé sur l'inégalité  $P'T \leq PT$  pour  $T \in \bar{D}$ .

Soit  $A_i \neq A_j$ . Le lieu géométrique des points équidistants de  $A_i$  et  $A_j$  est un plan qui divise l'espace en deux domaines, dont celui contenant  $A_i$  est appelé  $F_{ij}$ . Le produit  $C_i = D \prod_{j \neq i} F_{ij}$  est un domaine borné et convexe, contenu dans  $D$ ;  $D = \bar{D} - F$  (comme ci-dessus) et  $\bar{D}$  est l'ensemble dont parle le lemme e).

f) Soit  $(K, A)$  la solution du problème. Nous disons que  $K$  est la décomposition de  $Q$  fournie par  $E_i = C_i E$ ,  $E$  étant l'ensemble caractéristique de  $Q$  et  $C_i$  les ensembles dont parle le lemme e).

En effet, pour une décomposition quelconque  $E_i$  de  $E$  on obtient

$$S(K, A) = \iiint_E f(P) k^2(P) dv, \quad (dv = dx dy dz),$$

où  $k(P) = PA_i$  pour  $P \in E_i$  ( $i=1, 2, \dots, n$ ). Il est évident que l'on a toujours

$$k(P) \geq h(P) = \min(PA_1, PA_2, \dots, PA_n)$$

et que, par conséquent, la valeur minimum de  $S(K, A)$ , pour un  $A$  donné, est obtenue lorsque  $k(P) = h(P)$  presque partout; or, ce cas n'est réalisé que par la *décomposition spéciale*  $\{E_i\}$  définie par les plans équidistants.



Les lemmes b), d) et f) montrent que la solution du problème ne peut être obtenue qu'en réalisant les conditions que l'on ait exactement  $n$  points  $A_i$  distincts, qu'ils soient les centroïdes des corps partiels  $K_i$  et que ces corps soient séparés deux à deux par les plans équidistants relatifs à leurs centroïdes respectifs.

Nous allons établir maintenant l'existence d'une solution, en suivant une voie directe.

Soit  $m$  la limite inférieure de  $S(X, Y)$  pour tous les  $(X, Y)$  possibles; on aura, pour une suite convenable  $(X^r, Y^r)$ ,

$$(2) \quad \lim_{r \rightarrow \infty} S(X^r, Y^r) = m.$$

Cette relation subsiste — à cause du lemme d) — lorsqu'on remplace chaque point  $Y_i^r$  par le point qui lui est le plus proche parmi les points de  $\bar{D}$ ; on peut donc supposer que tous les  $Y_i^r$  dans (2) appartiennent à  $\bar{D}$ . Il s'ensuit que l'on peut extraire de  $\{Y^r\}$  une suite convergente vers  $Y^0$ : on entend par là  $\lim_{r \rightarrow \infty} Y_i^r = Y_i^0$  ( $i=1, 2, \dots, n$ ) pour la suite partielle (dont les termes sont désignés ici par les mêmes lettres que ceux de la suite primitive). On en tire

$$(3) \quad \lim_{r \rightarrow \infty} S(X^r, Y^0) = m$$

pour une certaine suite  $\{X^r\}$ . Or, il est possible que les  $n$  points  $Y_i^0$  ne soient pas tous distincts; soit par exemple  $Y_1^0 = Y_2^0$ . On aura alors

$$(4) \quad I(X_1^r, Y_1^0) + I(X_2^r, Y_2^0) = I(X_1^r + X_2^r, Y_1^0).$$

$X_1^r + X_2^r$  est un nouveau corps partiel qui remplace dans  $S$  deux corps de la division primitive; ce changement n'altère pas la valeur de  $S$  à cause de (4). En poursuivant cette réduction de la somme  $S$  on aboutit à une expression  $(X^r, Y^0)$  avec  $s$  termes  $Y_i^0$  distincts ( $i=1, 2, \dots, s \leq n$ ) et la formule (3) subsiste pour l'expression nouvelle. Construisons maintenant la division spéciale relative à  $Y_i^0$  ( $i=1, 2, \dots, s$ ) (cf. lemme f) (cette division se réduit à  $X_1^0 = 0$  dans le cas exceptionnel  $s=1$ ). Cette division  $X^0$  ne dépend pas de  $r$  et on obtient

$$(5) \quad S(X^0, Y^0) \geq S(X^r, Y^0).$$

(3) et (5) donnent

$$(6) \quad S(X^0, Y^0) = m.$$

Le nombre des corps partiels dans  $X^0$  est  $s \leq n$ ; or, l'inégalité  $s < n$  est incompatible avec (6) d'après le lemme d); on voit donc que  $s = n$  ( $X^0, Y^0$ ) est bien la solution du problème principal de cette Note. Ce résultat démontre la possibilité de satisfaire simultanément aux conditions nécessaires, formulées à la fin du paragraphe 1. Pour un corps  $Q$  donné, le minimum  $m$  est une fonction d'une seule variable  $n$ ; l'égalité  $s = n$

montre que cette fonction est décroissante; il est aussi facile à voir que l'on a  $\lim_{n \rightarrow \infty} m(n) = 0$ .

La solution du problème,  $Q$  et  $n$  étant donnés, est en général unique, mais il y a des corps avec une infinité de solutions: la sphère homogène en est un pour tous les  $n$ . Bien entendu, il y a des corps avec un nombre fini de solutions différentes: un cube homogène, par exemple, en a trois pour  $n=2$ .

Même s'il n'y a qu'une seule solution, les conditions nécessaires du paragraphe 1 ne sont pas suffisantes: un bloc homogène  $|x| \leq a$ ,  $|y| \leq b$ ,  $|z| \leq c$  avec  $a > b > c$  fournit un exemple pour  $n=2$ ; en effet, en le coupant par le plan  $x=0$  en deux et en posant  $A_1 = (-a/2, 0, 0)$ ,  $A_2 = (a/2, 0, 0)$ , on obtient la solution unique, mais on peut couper le bloc par le plan  $y=0$  ou par le plan  $x=0$  pour obtenir deux autres divisions qui satisfont aux conditions nécessaires sans toutefois fournir le minimum.

Dans les applications il n'y aura pour la plupart qu'un seul  $(K, A)$  satisfaisant aux conditions nécessaires; il paraît que dans ce cas la solution peut être trouvée en partant d'une décomposition  $K^1$  arbitraire en  $n$  parties, en définissant  $A'_i$  ( $i=1, 2, \dots, n$ ) comme centroïde de  $K'_i$ , et  $K^{r+1}$  comme la décomposition spéciale relative à  $A^r$ , pour tous les  $r$  naturels; or, il nous manque une démonstration de la convergence de  $(K^r, A^r)$  pour  $r \rightarrow \infty$  vers la solution cherchée.

La question se pose, si l'existence d'une infinité de solutions pour chaque  $n (\geq 2)$  implique que  $Q$  soit un corps de révolution. Une autre question: si l'on a pour un  $Q$  donné plusieurs solutions pour certains  $n$ , l'ensemble de ces  $n$  est-il nécessairement infini?

Diverses questions, par exemple celles des types en anthropologie, ou bien d'autres, d'ordre pratique, comme celles de la normalisation des objets industriels, exigent pour leur solution la détermination de  $n$  représentants fictifs d'une nombreuse population, choisis de manière à réduire autant que possible les écarts entre les éléments de la population et ceux de l'échantillon, l'écart étant mesuré entre tout élément réel et l'élément fictif qui lui est le plus proche — voilà les considérations qui ont conduit l'auteur à étudier le cas spécial traité dans cette Note.

## The Calculation of the Matrix Components of Energy for Hexagonal Close-Packed Structure

by

M. MIASEK

*Presented by L. INFELD on September 28, 1956*

The tight binding method originally proposed by Bloch [1] and worked out in detail by Slater and Koster [2], was used here to investigate the hexagonal close-packed structure.

It is known that this method in the approximation proposed in [2] gives satisfactory quantitative results, if applied to metals in which the electrons responsible for conduction are tightly bound. In particular, it shows correctly all the symmetry properties in energy bands.

Such metals are the transition metals Fe, Co and Ni, which have the  $3d$  shell partly and the  $4s$  shell completely filled. Fletcher and Wohlfarth [3] and Fletcher [4] applied the tight binding method to  $\beta$ -Ni, which has a face-centered cubic structure.

There is, however, a form of  $\alpha$ -Co and  $\alpha$ -Ni, which has a hexagonal close-packed structure, and the calculations reported below were performed in order to use them later to study the properties of the energy bands of these metals.

The matrix components of energy between the Bloch sums were calculated using first the general energy integrals ( $E$ -integrals) and subsequently the two-centre integrals. The general formula for the matrix component of energy is given by

$$H_{mn}^{\omega\omega'} = (m/n)_{\omega\omega'} = \exp[i\mathbf{k}(\mathbf{t}_\omega - \mathbf{t}_{\omega'})] \sum_i \exp(i\mathbf{k}\mathbf{r}_i) \int \varphi_m^*(\mathbf{r} - \mathbf{t}_{\omega'}) H \varphi_n(\mathbf{r} - \mathbf{r}_i - \mathbf{t}_\omega) d\mathbf{r},$$

where  $m$  and  $n$  denote the electronic states; in our case they are the  $s$  and  $d$  states.  $\mathbf{t}_\omega$  denotes the basis vectors.  $\varphi_m(\mathbf{r})$  is the eigenfunction of the electrons in an isolated atom. Energy integrals ( $E$ -integrals) have the form

$$E_{mn}(\mathbf{t}_{\omega'}, \mathbf{r}_i + \mathbf{r}_\omega) = \int \varphi_m^*(\mathbf{r} - \mathbf{t}_{\omega'}) H \varphi_n(\mathbf{r} - \mathbf{r}_i - \mathbf{t}_\omega) d\mathbf{r}.$$



The hexagonal close-packed lattice can be regarded as the simple hexagonal lattice with two atoms in the unit cell. The translation vectors of hexagonal lattice in a rectangular co-ordinate system are:  $\mathbf{A}_1 = (a/2, -\sqrt{3}a/2, 0)$ ,  $\mathbf{A}_2 = (a/2, \sqrt{3}a/2, 0)$ ,  $\mathbf{A}_3 = (0, 0, c)$ . The minimum distance in simple hexagonal lattice is  $a$  (lattice constant). Basis vectors are given by  $\mathbf{t}_1 = (0, 0, 0)$ ,  $\mathbf{t}_2 = (a/2, \sqrt{3}a/6, c/2)$ . The lattice obtained by starting from the central atom in the position  $(0, 0, 0)$  is denoted by the index 1, and the lattice obtained by a translation in the position  $(a/2, \sqrt{3}a/6, c/2)$  is denoted by 2. In the calculations we used a model of the hexagonal ideal close-packed lattice, i. e. we assumed  $|\mathbf{t}_2| = a$ . In this case  $c/a = 1.633$ . The assumption  $|\mathbf{t}_2| = a$  is satisfactory for  $\alpha$ -Co and  $\alpha$ -Ni, because for Co the ratio  $c/a$  has the value 1.62 and for Ni  $c/a = 1.64$ . It is now convenient to use different length units in different directions of the axes of rectangular co-ordinate systems. So we take  $a = a/2$ ,  $\beta = \sqrt{3}a/2$ ,  $\gamma = c/2$  in the  $x, y, z$  directions respectively.

In the formula for  $(m/n)_{\omega\omega'}$  we considered only the integrals representing the interaction between nearest neighbours. Regarding an atom in the position  $(0, 0, 0)$  we have six neighbours of the lattice 1, which are all in the same plane together with the central atom. Their positions in our rectangular co-ordinate system are  $(1, -1, 0)$ ,  $(2, 0, 0)$ ,  $(1, 1, 0)$ ,  $(-1, 1, 0)$ ,  $(-2, 0, 0)$ ,  $(-1, -1, 0)$ . The matrix components of energy are of the type  $(m/n)_{11}$ . We also have six neighbours belonging to the lattice 2, which are placed at a distance  $c/2$  above and below this plane. Their positions are  $(1, 1/3, 1)$ ,  $(-1, 1/3, 1)$ ,  $(0, -2/3, 1)$ ,  $(1, 1/3, -1)$ ,  $(-1, 1/3, -1)$ ,  $(0, -2/3, -1)$ . The matrix components of energy are here of the  $(m/n)_{12}$  type. Accordingly, the sum for  $(m/n)_{\omega\omega'}$  will contain six terms.  $(m/n)_{11}$  also contains the term with  $\mathbf{r}_i = (0, 0, 0)$ .

Since we consider six states (five  $d$  and one  $s$  states) and in the unit cell we have two atoms, the general matrix of energy has  $12 \times 12 = 144$  elements. Using Hermitian and symmetry properties of the Hamiltonian and the symmetry properties of the eigenfunctions of the isolated atoms we find that the matrix of energy has only 42 different components. Other components are either equal or conjugate to these. The symmetry properties of the Hamiltonian and of the eigenfunctions make it possible to reduce still further the number of different  $E$ -integrals which occur in the matrix components of energy.

In Table I we have given the matrix components of energy in terms of  $E$ -integrals. The abbreviation  $\xi = ak_x/2$ ,  $\eta = \sqrt{3}ak_y/2$ ,  $\zeta = ck_z/2$  has been used in this table. Since in many matrix components we have the same dependence on  $\xi, \eta, \zeta$ , we have written only one element of this type and on the right hand side of the table the components with the same dependence on  $\mathbf{k}$  are given. Evidently they differ only in the form of  $E$ -integrals.

Table I

$(s/s)_{11}$	$(s/3z^2 - r^2)_{11}$
$E_{s,s}(0, 0, 0) + 2E_{s,s}(1, -1, 0)(2 \cos \xi \cos \eta + \cos 2\xi)$	$(3z^2 - r^2/3z^2 - r^2)_{11}$
$(s/s)_{12}$	$(s/3z^2 - r^2)_{12}$
$2E_{s,s}(1, \frac{1}{3}, 1) \cos \zeta [(2 \cos \xi \cos \eta/3 + \cos 2\eta/3) +$ $+ i(2 \cos \xi \sin \eta/3 - \sin 2\eta/3)]$	$(3z^2 - r^2/3z^2 - r^2)_{12}$
$(s/xy)_{11}$	$(yz/xz)_{11}$
$2[E_{s,xy}(1, -1, 0) - E_{s,xy}(1, 1, 0)] \sin \xi \sin \eta +$ $+ 2i\{[E_{s,xy}(1, -1, 0) + E_{s,xy}(1, 1, 0)] \sin \xi \cos \eta +$ $+ E_{s,xy}(2, 0, 0) \sin 2\xi\}$	$(xy/x^2 - y^2)_{11}$ $(xy/3z^2 - r^2)_{11}$
$(s/xy)_{12}$	$(yz/xz)_{12}$
$-4E_{s,xy}(1, \frac{1}{3}, 1) \sin \xi \cos \zeta (\sin \eta/3 - i \cos \eta/3)$	$(xy/x^2 - y^2)_{12}$ $(xy/3z^2 - r^2)_{12}$
$(s/yz)_{11} = (s/xz)_{11} = (xy/yz)_{11} = (xy/xz)_{11} = (yz/x^2 - y^2)_{11} =$ $= (yz/3z^2 - r^2)_{11} = (xz/x^2 - y^2)_{11} = (xz/3z^2 - r^2)_{11} = 0$	
$(s/yz)_{12}$	$(xy/xz)_{12}$
$2 \sin \zeta [-2E_{s,yz}(1, \frac{1}{3}, 1) \cos \xi \sin \eta/3 +$ $+ E_{s,yz}(0, -\frac{2}{3}, 1) \sin 2\eta/3] + 2i \sin \zeta [2E_{s,yz}(1, \frac{1}{3}, 1) \cos \xi \cos \eta/3 +$ $+ E_{s,yz}(0, -\frac{2}{3}, 1) \cos 2\eta/3]$	$(yz/3z^2 - r^2)_{12}$ $(yz/x^2 - y^2)_{12}$
$(s/xz)_{12}$	$(xy/yz)_{12}$
$-4E_{s,xz}(1, \frac{1}{3}, 1) \sin \xi \sin \zeta (\cos \eta/3 + i \sin \eta/3)$	$(xz/x^2 - y^2)_{12}$ $(xz/3z^2 - r^2)_{12}$
$(s/x^2 - y^2)_{11}$	$(x^2 - y^2/3z^2 - r^2)_{11}$
$E_{s,x^2-y^2}(0, 0, 0) + 2[E_{s,x^2-y^2}(1, -1, 0) + E_{s,x^2-y^2}(1, 1, 0)] \cos \xi \cos \eta +$ $+ 2E_{s,x^2-y^2}(2, 0, 0) \cos 2\xi + 2i[E_{s,x^2-y^2}(1, 1, 0) -$ $- E_{s,x^2-y^2}(1, -1, 0)] \cos \xi \sin \eta$	
$(s/x^2 - y^2)_{12}$	$(xy/xy)_{12}$
$2 \cos \zeta [2E_{s,x^2-y^2}(1, \frac{1}{3}, 1) \cos \xi \cos \eta/3 +$ $+ E_{s,x^2-y^2}(0, \frac{2}{3}, 1) \cos 2\eta/3] +$ $+ 2i \cos \zeta [2E_{s,x^2-y^2}(1, \frac{1}{3}, 1) \cos \xi \sin \eta/3 -$ $- E_{s,x^2-y^2}(0, -\frac{2}{3}, 1) \sin 2\eta/3]$	$(yz/yz)_{12}$ $(xz/xz)_{12}$ $(x^2 - y^2/x^2 - y^2)_{12}$ $(x^2 - y^2/3z^2 - r^2)_{12}$
$(xy/xy)_{11}$	$(yz/yz)_{11}$
$E_{xy,xy}(0, 0, 0) + 4E_{xy,xy}(1, -1, 0) \cos \xi \cos \eta +$ $+ 2E_{xy,xy}(2, 0, 0) \cos 2\xi$	$(xz/xz)_{11}$ $(x^2 - y^2/x^2 - y^2)_{11}$

We have here 72 different  $E$ -integrals, the calculation of which would be very laborious. Therefore it is convenient to use the two-centre approximation. To express the  $E$ -integrals in terms of the two-centre integrals ( $ss\sigma$ ), ( $sd\sigma$ ), ( $dd\sigma$ ), ( $dd\pi$ ), ( $dd\delta$ ) we used Table I in paper [2].

The matrix components expressed in terms of the two-centre integrals are given in Table II.

Table II

$(s/s)_{11}$	$s_0 + 2(ss\sigma)_1(2\cos\xi\cos\eta + \cos 2\xi)$
$(s/s)_{12}$	$2(ss\sigma)_1\zeta\cos\zeta[(2\cos\xi\cos\eta/3 + \cos 2\eta/3) + i(2\cos\xi\sin\eta/3 - \sin 2\eta/3)]$
$(s/xy)_{11}$	$-3(s\bar{d}\sigma)_1\sin\xi\sin\eta$
$(s/xy)_{12}$	$-(s\bar{d}\sigma)_1\sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$
$(s/yz)_{11}$	0
$(s/yz)_{12}$	$-2\sqrt{\frac{2}{3}}(s\bar{d}\sigma)_1\sin\zeta[(\cos\xi\sin\eta/3 + \sin 2\eta/3) - i(\cos\xi\cos\eta/3 - \cos 2\eta/3)]$
$(s/xz)_{11}$	0
$(s/xz)_{12}$	$-2\sqrt{2}(s\bar{d}\sigma)_1\sin\xi\sin\zeta(\cos\eta/3 + i\sin\eta/3)$
$(s/x^2 - y^2)_{11}$	$-\sqrt{3}(s\bar{d}\sigma)_1(\cos\xi\cos\eta - \cos 2\xi)$
$(s/x^2 - y^2)_{12}$	$\frac{1}{3}\sqrt{3}(s\bar{d}\sigma)_1\cos\zeta[(\cos\xi\cos\eta/3 - \cos 2\eta/3) + i(\cos\xi\sin\eta/3 + \sin 2\eta/3)]$
$(s/3z^2 - r^2)_{11}$	$-(s\bar{d}\sigma)_1(2\cos\xi\cos\eta + \cos 2\xi)$
$(s/3z^2 - r^2)_{12}$	$(s\bar{d}\sigma)_1\cos\zeta[(2\cos\xi\cos\eta/3 + \cos 2\eta/3) + i(2\cos\xi\sin\eta/3 - \sin 2\eta/3)]$
$(xy/xy)_{11}$	$d_0 + 2(dd\pi)_1\cos 2\xi + [\frac{9}{4}(dd\sigma)_1 + (dd\pi)_1 + \frac{3}{4}(dd\delta)_1]\cos\xi\cos\eta$
$(xy/xy)_{12}$	$2\cos\zeta\{\frac{1}{3}[(dd\pi)_1 + 2(dd\delta)_1]\cos 2\eta/3 + \frac{1}{2}[\frac{1}{4}(dd\sigma)_1 + (dd\pi)_1 + \frac{1}{4}(dd\delta)_1]\cos\xi\cos\eta/3\} + 2i\cos\zeta\{-\frac{1}{3}[(dd\pi)_1 + 2(dd\delta)_1]\sin 2\eta/3 + \frac{1}{2}[\frac{1}{4}(dd\sigma)_1 + (dd\pi)_1 + \frac{1}{4}(dd\delta)_1]\cos\xi\sin\eta/3\}$
$(xy/xy)_{11}$	0
$(xy/yz)_{12}$	$-\sqrt{\frac{2}{3}}[\frac{1}{2}(dd\sigma)_1 + \frac{4}{3}(dd\pi)_1 - \frac{1}{6}(dd\delta)_1] \times \sin\xi\sin\zeta(\cos\eta/3 + i\sin\eta/3)$
$(xy/xz)_{11}$	0
$(xy/xz)_{12}$	$2\sqrt{2}\sin\zeta\{\frac{1}{4}[-(dd\sigma)_1 + (dd\delta)_1]\cos\xi\sin\eta/3 + \frac{1}{3}[-(dd\pi)_1 + (dd\delta)_1]\sin 2\eta/3\} + 2\sqrt{2}i\sin\zeta\{\frac{1}{4}[(dd\sigma)_1 - (dd\delta)_1]\cos\xi\cos\eta/3 + \frac{1}{3}[-(dd\pi)_1 + (dd\delta)_1]\cos 2\eta/3\}$
$(xy/x^2 - y^2)_{11}$	$\sqrt{3}[\frac{3}{4}(dd\sigma)_1 - (dd\pi)_1 + \frac{1}{4}(dd\delta)_1]\sin\xi\sin\eta - \frac{1}{3}\sqrt{3}[\frac{1}{4}(dd\sigma)_1 - \frac{1}{3}(dd\pi)_1 + \frac{1}{12}(dd\delta)_1] \times \sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$
$(xy/x^2 - y^2)_{12}$	$-\frac{1}{3}\sqrt{3}[\frac{1}{4}(dd\sigma)_1 - \frac{1}{3}(dd\pi)_1 + \frac{1}{12}(dd\delta)_1] \times \sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$
$(xy/3z^2 - r^2)_{11}$	$\frac{3}{2}[(dd\sigma)_1 - (dd\delta)_1]\sin\xi\sin\eta - [\frac{1}{2}(dd\sigma)_1 - \frac{4}{3}(dd\pi)_1 + \frac{5}{6}(dd\delta)_1] \times \sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$
$(xy/3z^2 - r^2)_{12}$	$-\frac{1}{2}(dd\sigma)_1 - \frac{4}{3}(dd\pi)_1 + \frac{5}{6}(dd\delta)_1] \times \sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$
$(yz/yz)_{11}$	$d_0 + [3(dd\pi)_1 + (dd\delta)_1]\cos\xi\cos\eta + 2(dd\delta)_1\cos 2\xi$
$(yz/yz)_{12}$	$\frac{2}{3}\cos\zeta\{[2(dd\sigma)_1 + \frac{1}{3}(dd\pi)_1 + \frac{2}{3}(dd\delta)_1]\cos 2\eta/3 + [(dd\sigma)_1 + \frac{1}{6}(dd\pi)_1 + \frac{1}{6}(dd\delta)_1]\cos\xi\cos\eta/3\} + \frac{2}{3}i\cos\zeta\{-[2(dd\sigma)_1 + \frac{1}{3}(dd\pi)_1 + \frac{2}{3}(dd\delta)_1]\sin 2\eta/3 + [(dd\sigma)_1 + \frac{1}{6}(dd\pi)_1 + \frac{1}{6}(dd\delta)_1]\cos\xi\sin\eta/3\}$
$(yz/xz)_{11}$	$\sqrt{3}[-(dd\pi)_1 + (dd\delta)_1]\sin\xi\sin\eta$
$(yz/xz)_{12}$	$-\frac{2}{3}\sqrt{3}[(dd\sigma)_1 - \frac{5}{6}(dd\pi)_1 - \frac{1}{6}(dd\delta)_1] \times \sin\xi\cos\zeta(\sin\eta/3 - i\cos\eta/3)$



$$\begin{aligned}
(yz/x^2 - y^2)_{11} & 0 \\
(yz/x^2 - y^2)_{12} & \frac{2}{3} \sqrt{2} \sin \zeta \left\{ \left[ \frac{1}{2} (dd\sigma)_1 + \frac{1}{3} (dd\pi)_1 - \frac{5}{6} (dd\delta)_1 \right] \sin 2\eta/3 + \right. \\
& \quad \left. + \left[ -\frac{1}{4} (dd\sigma)_1 + \frac{4}{3} (dd\pi)_1 - \frac{1}{2} (dd\delta)_1 \right] \cos \xi \sin \eta/3 \right\} + \\
& \quad + \frac{2}{3} \sqrt{2} i \sin \zeta \left\{ \left[ \frac{1}{2} (dd\sigma)_1 + \frac{1}{3} (dd\pi)_1 - \frac{5}{6} (dd\delta)_1 \right] \cos 2\eta/3 + \right. \\
& \quad \left. + \left[ \frac{1}{4} (dd\sigma)_1 - \frac{4}{3} (dd\pi)_1 + \frac{1}{2} (dd\delta)_1 \right] \cos \xi \cos \eta/3 \right\} \\
(yz/3z^2 - r^2)_{11} & 0 \\
(yz/3z^2 - r^2)_{12} & -\frac{2}{3} \sqrt{2} \left[ \frac{1}{2} (dd\sigma)_1 - \frac{1}{3} (dd\pi)_1 - \frac{1}{6} (dd\delta)_1 \right] \sin \zeta \times \\
& \quad \times [(\cos \xi \sin \eta/3 + \sin 2\eta/3) - i(\cos \xi \cos \eta/3 - \cos 2\eta/3)] \\
(xz/xz)_{11} & d_0 + [(dd\pi)_1 + 3(dd\delta)_1] \cos \xi \cos \eta + 2(dd\pi)_1 \cos 2\xi \\
(xz/xz)_{12} & 2 \cos \zeta \left\{ \frac{1}{3} [2(dd\pi)_1 + (dd\delta)_1] \cos 2\eta/3 + \right. \\
& \quad \left. + [(dd\sigma)_1 + \frac{1}{2} (dd\pi)_1 + \frac{1}{2} (dd\delta)_1] \cos \xi \cos \eta/3 \right\} + \\
& \quad + 2i \cos \zeta \left\{ -\frac{1}{3} [2(dd\pi)_1 + (dd\delta)_1] \sin 2\eta/3 + \right. \\
& \quad \left. + [(dd\sigma)_1 + \frac{1}{2} (dd\pi)_1 + \frac{1}{2} (dd\delta)_1] \cos \xi \sin \eta/3 \right\} \\
(xz/x^2 - y^2)_{11} & 0 \\
(xz/x^2 - y^2)_{12} & -\sqrt{\frac{2}{3}} \left[ \frac{1}{2} (dd\sigma)_1 + \frac{4}{3} (dd\pi)_1 - \frac{1}{6} (dd\delta)_1 \right] \times \\
& \quad \times \sin \xi \sin \zeta (\cos \eta/3 + i \sin \eta/3) \\
(xz/3z^2 - r^2)_{11} & 0 \\
(xz/3z^2 - r^2)_{12} & -\sqrt{2} [(dd\sigma)_1 - \frac{2}{3} (dd\pi)_1 - \frac{1}{3} (dd\delta)_1] \times \\
& \quad \times \sin \xi \sin \zeta (\cos \eta/3 + i \sin \eta/3) \\
(x^2 - y^2/x^2 - y^2)_{11} & d_0 + [\frac{3}{4} (dd\sigma)_1 + 3(dd\pi)_1 + \frac{1}{4} (dd\delta)_1] \cos \xi \cos \eta + \\
& \quad + \frac{1}{2} [3(dd\sigma)_1 + (dd\delta)_1] \cos 2\xi \\
(x^2 - y^2/x^2 - y^2)_{12} & \frac{2}{3} \cos \zeta \left\{ \left[ \frac{1}{4} (dd\sigma)_1 + \frac{2}{3} (dd\pi)_1 + \frac{2}{3} (dd\delta)_1 \right] \cos 2\eta/3 + \right. \\
& \quad \left. + \frac{1}{2} \left[ \frac{1}{4} (dd\sigma)_1 + \frac{1}{3} (dd\pi)_1 + \frac{2}{3} (dd\delta)_1 \right] \cos \xi \cos \eta/3 \right\} + \\
& \quad + \frac{2}{3} i \cos \zeta \left\{ -\left[ \frac{1}{4} (dd\sigma)_1 + \frac{2}{3} (dd\pi)_1 + \frac{2}{3} (dd\delta)_1 \right] \sin 2\eta/3 + \right. \\
& \quad \left. + \frac{1}{2} \left[ \frac{1}{4} (dd\sigma)_1 + \frac{1}{3} (dd\pi)_1 + \frac{2}{3} (dd\delta)_1 \right] \cos \xi \sin \eta/3 \right\} \\
(x^2 - y^2/3z^2 - r^2)_{11} & \frac{1}{2} \sqrt{3} [(dd\sigma)_1 - (dd\delta)_1] (\cos \xi \cos \eta - \cos 2\xi) \\
(x^2 - y^2/3z^2 - r^2)_{12} & \frac{2}{3} \sqrt{3} \cos \zeta \left\{ \left[ -\frac{1}{4} (dd\sigma)_1 + \frac{2}{3} (dd\pi)_1 - \frac{5}{12} (dd\delta)_1 \right] \cos 2\eta/3 + \right. \\
& \quad \left. + \frac{1}{4} \left[ (dd\sigma)_1 - \frac{2}{3} (dd\pi)_1 + \frac{5}{12} (dd\delta)_1 \right] \cos \xi \cos \eta/3 \right\} + \\
& \quad + \frac{2}{3} \sqrt{3} i \cos \zeta \left\{ -\left[ -\frac{1}{4} (dd\sigma)_1 + \frac{2}{3} (dd\pi)_1 - \frac{5}{12} (dd\delta)_1 \right] \sin 2\eta/3 + \right. \\
& \quad \left. + \left[ \frac{1}{4} (dd\sigma)_1 - \frac{2}{3} (dd\pi)_1 + \frac{5}{12} (dd\delta)_1 \right] \cos \xi \sin \eta/3 \right\} \\
(3z^2 - r^2/3z^2 - r^2)_{11} & d_0 + \frac{1}{2} [(dd\sigma)_1 + 3(dd\delta)_1] (2 \cos \xi \cos \eta + \cos 2\xi) \\
(3z^2 - r^2/3z^2 - r^2)_{12} & \left[ \frac{1}{2} (dd\sigma)_1 + \frac{4}{3} (dd\pi)_1 + \frac{1}{6} (dd\delta)_1 \right] \cos \xi \times \\
& \quad \times [(2 \cos \xi \cos \eta/3 + \cos 2\eta/3) + i(2 \cos \xi \sin \eta/3 - \sin 2\eta/3)]
\end{aligned}$$

It may be seen from the above that the two-centre approximation considerably simplifies the calculations. Instead of the 72  $E$ -integrals we now have only 7 integrals,  $s_0$ ,  $d_0$ ,  $(ss\sigma)_1$ ,  $(s\delta\sigma)_1$ ,  $(dd\sigma)_1$ ,  $(dd\pi)_1$  and  $(dd\delta)_1$ . When we know the  $\mathbf{k}$  dependence of the matrix components and

the values of the two-centre integrals, the symmetry properties of the energy bands can be studied.

I wish to thank Dr M. Suffczyński for his great interest in this work and for valuable discussions.

INSTITUTE OF PHYSICS, POLISH ACADEMY OF SCIENCES

#### REFERENCES

- [1] F. Bloch, Zft. Phys. **52**, (1928), 555.
- [2] J. C. Slater, Koster G. F., Phys. Rev. **94**, (1954), 1498.
- [3] G. C. Fletcher, E. P. Wohlfarth, Phil. Mag. **42**, (1951), 106.
- [4] G. C. Fletcher, Proc. Phys. Soc. **65A**, (1952), 192.

# Investigation of an Electromagnetic Cascade of Very High Energy in the First Stage of Its Development in Nuclear Emulsions

by

M. MIEŚOWICZ, O. STANISZ and W. WOLTER

*Presented by H. NIEWODNICZAŃSKI on September 18, 1956*

Information on some examples of high energy showers of electron pairs has recently been published. The development of some of these showers seems to show that they may be initiated by several photons [1]-[5]. At the same time the range of energy, in which electron showers initiated by a single photon have been found and investigated in emulsions, has been extended to above  $10^{11}$  eV [4], [6], [7].

One of the anomalies found in high energy showers is the deficiency of low energy pairs as compared with the number of high energy pairs. This effect has been reported for showers assumed to be initiated by several photons by Debenedetti et al. [2], and for a shower probably initiated by a single photon, by the authors of the present paper [6]. It was suggested by Wataghin [8] that these anomalies have a general character for all electronic showers of high energy ( $\simeq 10^{12}$  eV). Dyson [9] suggested that this anomaly might be explained on the basis of the theories of Ter Mikaelian [10] and Landau and Pomerančuk [11]. It is very important in this connection to determine whether the anomaly really exists in the showers initiated by a single photon of very high energy. It therefore seems very desirable to carry out investigations on cascades deriving from single photons of the highest energies at their beginning. At greater depths these anomalies should disappear because of the decrease in the energy of the electrons of the cascade.

Hitherto there were known in the literature only two cascades, generated by single photons of energies of  $\simeq 10^{12}$  eV. These were recently analysed by Pinkau [7].

The present paper is intended to give an account of an analysis of a cascade generated probably by a single photon of an energy approaching the value of  $10^{12}$  eV, particular consideration being given to the beginning



of the development of the cascade where we could expect to find such an anomaly. Preliminary results of the measurements of this cascade have already been published [6].

The shower consists of high energy pairs, 14 of which are generated within 1.55 rad. length from the origin of the first pair. Most of these pairs are included within a very narrow cone of an angle of about  $10^{-4}$  radians. The length of the cascade in one emulsion is about 4.2 mm. The data on the observed pairs are given in Table I. Most of the pairs are apparent tridents, as can be seen from column 7 of Table I.

The evaluation of energy of the primary photon initiating the cascade was carried out with the energy spectrum of electrons at a depth of 2.5 rad. length. The scattering measurements were made on the tracks contained within a circle of a radius of  $250 \mu$  from the core. The number of tracks of electrons with energies higher than  $5 \cdot 10^8$  eV within the circle of a radius of  $200 \mu$  was 34. The calculations of the full number of electrons in this energy interval at this depth, on the basis of the lateral distribution of particles, were carried out by a method described by Pinkau [7]. Then, using the tables of Janossy and Messel [12], the energy  $E_0$  of the primary photon could be found if we assumed the equipartition of energy for particles of the first pair. From these measurements we obtained  $E_0 = (7.0^{+3.4}_{-2.6}) \times 10^{11}$  eV. The limits of error of the value of  $E_0$  were found, taking Janossy's [13] standard deviations for a given number of tracks and finding the values of energy corresponding to the extreme number of electrons of energies greater than  $5 \cdot 10^8$  eV. From the value of  $E_0$  the numbers of electrons of different energies for various radii were obtained, and they are in agreement with the observed values within Janossy's standard deviations.

We obtained information on the energy of the primary pair independently in another way, from measurements of ionization on the first part of the unresolved track of the first pair. From these measurements [14]-[16] it was possible with the use of the formula given by Čudakov [14] to estimate the opening angle of the first pair. We obtained the value  $\omega = 3 \cdot 10^{-6}$  radians, and hence from the formula  $E = 4mc^2/\omega$  we obtained for  $E_0$  the value  $\sim 10^{12}$  eV. We consider these measurements as only approximate, since in the formula from which we evaluated the angle there are constants which are not accurately known.

We have now investigated the spectrum of electrons at a depth of 1.55 rad. length. In Fig. 1 the integral energy spectrum of electrons observed at this depth is shown. This spectrum is obtained directly from the figures given in Table I, taking into account the energy losses of electrons by "bremsstrahlung" and ionization. The circles show the numbers of electrons with energy greater than the given value. The vertical lines give the uncertainty of energy determination. The full line shows the spectrum of electrons calculated from the cascade theory (Approximate).

mation B) [12] on the assumption that the energy of the primary photon is  $E_0 = 7.10^{11}$  eV as mentioned above. The dotted lines represent the spectra calculated for energies giving the limits of error of the primary photon energy ( $4.4$  or  $10.4 \times 10^{11}$  eV).

As may be seen from Fig. 1 there is a deficiency in the number of electrons of energy lower than  $5.10^8$  eV as compared with the prediction of the cascade theory.

Qualitatively it is possible to explain this effect according to the suggestions of Dyson [9] on the basis of the theory of Ter-Mikaelian [10] and Landau and Pomerančuk [11]. According to these theories, if an




Fig. 1.

electron moves through photographic emulsion with an energy  $E$ , greater than  $\simeq 10^{10}$  eV, the spectrum of "bremsstrahlung" photons radiated by this electron, for photon energies lower than a critical energy  $k_c$ , is changed from the Bethe-Heitler form  $\left(\sim \frac{dk}{k}\right)$  to the Landau-Pomerančuk form  $\left(\sim \frac{dk}{\sqrt{k}}\right)$ , and is strongly cut down for low photon energies. For photographic emulsions,  $k_c \simeq 8 \cdot 10^{-15} E^2$ . This means that for the two electrons of our first pair the energy of photons below which we might expect a lack of photons is  $k_c \simeq 10^9$  eV. Since at the depth of  $t = 1.55$  rad. length the observed energy spectrum of electrons is strongly influenced by the spectrum of the photons radiated by the first pair, it is possible to explain the observed deficiency of electrons of energies  $10^8$  eV and lower by the lack of photons of these energies in the radiation emitted by the first two electrons. This explanation is supported by direct observation of pairs generated on the tracks of the first pair. It can be seen (Table I) that no photons of energy smaller than  $10^9$  eV were converted on the left track, whereas some pairs with energies  $\sim 10^8$  are seen

only on the right track. This might be explained by assuming an asymmetry of energies in the two electrons of the first pair. In consequence of the  $k_c \sim E^2$  dependence,  $k_c$  may differ for both electrons of the first pair by one order of magnitude.

It seems that the observed deficiency of low energy electrons at the depth  $t=1.55$  rad. length is outside the limits of the probable fluctuations. Nevertheless it is not possible to conclude quantitatively from this example whether it is in accordance with the theory. It is thus very desirable to check this effect in other high energy cascades generated by single photons of energy  $10^{12}$  eV or higher.

TABLE I

Pair	Distance from the point of origin of the first pair $\mu$	$E_1$ eV	$E_2$ eV	$E_1 + E_2$ eV	Most probable parent pair	Radial distance from the nearest electron track $\mu$
1	0	—	—	$7.0 \times 10^{11}$	—	—
2	9400	$2.0 \times 10^8$	$1.0 \times 10^8$	$3.0 \times 10^8$	1(l) or 1(r) *)	$< 0.5_0$
3	16400	$1.3 \times 10^8$	$1.0 \times 10^7$	$1.4 \times 10^8$	1(r)	$< 0.4_5$
4	17500	$6.0 \times 10^8$	$\geq 7.0 \times 10^8$	$\geq 1.3 \times 10^9$	1(l)	$< 0.2_6$
5	23800	$1.3 \times 10^8$	$8.0 \times 10^7$	$2.1 \times 10^8$	1(r)	$1.0_8$
6	27800	$\geq 1.0 \times 10^9$	$7.5 \times 10^8$	$\geq 1.8 \times 10^9$	1(l)	$< 0.2_3$
7	30000	$< 6.0 \times 10^7$	$6.0 \times 10^7$	$< 1.2 \times 10^8$	?	12.6
8	30600	—	—	$> 1.3 \times 10^{10}$ (a)	1(l)	$< 0.4_5$
9	34900	—	—	$5.2 \times 10^9$ (b)	1(l) or 8	$< 0.2_3$
10	38700	$1.0 \times 10^7$	$6.8 \times 10^6$	$1.7 \times 10^7$	6	$< 0.4_5$
11	40900	—	—	$> 1.0 \times 10^{10}$ (c)	1(r)	$1.3_1$
12	41600	—	—	$> 3.9 \times 10^{10}$ (a)	1(l)	$< 0.8_4$
13	42800	—	—	$> 2.6 \times 10^{10}$ (a)	1(l) or 12	$< 1.3_3$
14	43500	$\geq 2.7 \times 10^8$	$1.0 \times 10^8$	$\geq 3.7 \times 10^8$	8	$0.9_5$

\*) — (l) or (r) means left or right track of the pair.

(a) — the lower limit of energy estimated from the opening angle of the core containing several tracks using the formula of Borsellino. It is highly probable that this value is underestimated, since the influence of multiple scattering is neglected.

(b) — from relative scattering measurements on the assumption of equipartition of energy.

(c) — as in (a) — measurements taken between the tracks of this pair.



The authors wish to express their thanks to Miss A. Jurak for the work done in the course of scanning for pairs and in the scattering measurements.

DEPARTMENT OF COSMIC RAYS, NUCLEAR RESEARCH INSTITUTE, POLISH ACADEMY OF SCIENCES, CRACOW

## REFERENCES

- [1] M. Schein, D. M. Haskin, R. G. Glasser, *Phys. Rev.* **95** (1954), 855.
- [2] A. Debenedetti, C. M. Garelli, L. Tallone, M. Vigone, G. Wataghin, *Nuovo Cimento* **12** (1954), 954; **2** (1955), 220; *ibid.* **3** (1956), 226.
- [3] A. Milone, *Pisa Conference*, June 1955.
- [4] M. Koshiba, M. F. Kaplon, *Phys. Rev.* **100** (1955), 327.
- [5] L. Barbanti Silva, C. Bonacini, C. Depietri, I. Iori, G. Lovera, R. Perilli Federi, A. Roveri, *Nuovo Cimento* **3** (1956), 1465.
- [6] M. Mięsowicz, W. Wolter, O. Stanisiz, *Pisa Conference*, June 1955; A. Jurak, M. Mięsowicz, O. Stanisiz, W. Wolter, *Bull. Acad. Polon. Sci., Cl. III*, **3** (1955), 369.
- [7] K. Pinkau, *Nuovo Cimento* **3** (1956), 1285.
- [8] G. Wataghin, *Proc. of the Sixth Rochester Conference IX*, **24** (1956).
- [9] F. I. Dyson, *Proc. of the Sixth Rochester Conference IX*, **29** (1956).
- [10] M. L. Ter-Mikaelian, *Izv. Akad. Nauk SSSR, Ser. fiz.* **19** (1955), 657.
- [11] L. D. Landau, I. Pomerančuk, *Dokl. Akad. Nauk SSSR* (1953), 92, 535, 735. (See also: E. L. Feinberg and I. Pomerančuk, *Supplemento, Nuovo Cimento* **3** (1956), 652).
- [12] L. Janossy, H. Messel, *Proc. Roy. Irish Acad.* **54** (A) (1951), 217.
- [13] L. Janossy, *Acta Phys. Hung.* **II** (1952), 289.
- [14] A. E. Čudakov, *Izv. Akad. Nauk SSSR Ser. fiz.* **14** (1955), 651.
- [15] D. H. Perkins, *Phil. Mag.* **46** (1955), 1146.
- [16] W. Wolter, M. Mięsowicz, *Nuovo Cimento* **4** (1956), 648.



## The Ratio of the Number of Photons to the Number of Electrons in Extensive Air Showers of Cosmic Radiation

by

J. BABECKI, L. JURKIEWICZ, J. M. MASSALSKI and M. MIĘSOWICZ

*Presented by H. NIEWODNICZAŃSKI on September 19, 1956*

Recently various authors have investigated the transition curve of extensive air shower particles in lead [1]-[4]. With these measurements is connected the problem of the ratio  $p/e$  of the number of photons to the number of electrons in extensive showers.

In the methods of measurement of  $p/e$ , in which we cover one or more counter trays with the absorber, we may evaluate from experimental data the quantity

$$(1) \quad R(t) = P_e(t) + \frac{p}{e} \cdot P_p(t),$$

where  $P_e$  and  $P_p$  are the probabilities that an electron or a photon respectively, falling on the absorber of a given thickness  $t$ , will produce beneath the absorber at least one secondary electron, which operates the counter tray. The functions  $P_e(t)$  and  $P_p(t)$  have been evaluated theoretically by Arley [5], and they are conventionally adopted by most authors (except Millar) considering the problem of the ratio  $p/e$  in spite of the objections against Arley's theory. Knowing the values of  $P_e$  and  $P_p$ , it is possible to evaluate the ratio  $p/e$  from formula (1).

In the anti-coincidence method we make use of the counter telescope, in which an absorber is placed between the two trays; then we estimate the rate of events in which only the lower tray of the telescope is operated. The rate of these anti-coincidences depends only on the expression  $p/e \cdot P_p$ , but does not depend on  $P_e$ , and this decides on the superiority of the anti-coincidence method.

In the first series of experiments concerning measurements of  $p/e$ , the values obtained were rather low, namely 0.25 [1] and 0.12 and 0.3 [2]. It was difficult to explain, from the point of view of the one-dimensional cascade theory, such a low value of  $p/e$ , since it predicts for this ratio



the value  $\geq 1$ . Milone in the last of his papers [3] gives for this ratio the value  $0.75 \pm 0.20$ , and Massalski [4], 0.8 and  $1.0 \pm 0.3$ .

### Apparatus

We used in our experiment a modified version of the apparatus previously employed. The detector of extensive showers was composed of three counter trays  $A, B, C$ , each of an area  $S \doteq 0.468$  sq. m., placed in the corners of an equilateral triangle with sides of 3.4 m. In the centre of the triangle was placed a telescope composed of three counter trays  $D, E, F$ , each of an area  $s = r \cdot S = 0.290$  sq. m. Tray  $E$  was placed immediately above tray  $F$ , whereas between trays  $D$  and  $E$  it was possible to place an absorber 15 cm. thick. The telescope was enclosed by lead walls 10 cm. thick and a lead base 5 cm. deep. All counters were made of brass tubes with walls 1 mm. thick. The measurements were performed in Cracow at an altitude of 229 m. above sea level.

### Principles of measurements and their results

Two series of measurements were performed. In the first series lead absorbers from 0 to 25 cm. thick were placed over the telescope  $DEF$ . In the second series lead absorbers 0 to 15 cm. thick were placed between the trays  $D$  and  $E$ . In each of these series the following were registered: threefold coincidences  $(ABC) = T$ ; fourfold coincidences  $(TD)$ ,  $(TE)$ ,  $(TF)$ ; fivefold coincidences  $(TDE)$ ,  $(TDF)$ ,  $(TEF)$ ; sixfold coincidences  $(TDEF)$ . In the anti-coincidence method we used an arrangement in which one tray ( $D$ ) was placed over the absorber, while the second tray ( $E$ ) was below it. The third tray of the telescope ( $F$ ) was added to eliminate the influence of the low energy penetrating photons of only several MeV which are present in relatively large numbers beneath thicker lead absorbers, and distinctly contribute to the registered rates of coincidences when below the absorber one counter tray is operated; they do not, however, influence the readings when two trays under the absorber are operated in coincidence [6], [7]. All the coincidences may be divided into two groups:  $A$  and  $B$ . Group  $A$  comprises the measurements of transition curves in which the absorber is placed over the telescope containing one, two or three trays of counters. The rates of coincidences of group  $A$  may be expressed by the formula

$$(2) \quad A = \int_0^{\infty} K x^{-(\gamma+1)} (1 - e^{-Sx})^3 (1 - e^{-(P_e + p/e \cdot P_p + k) r' r \cdot Sx}) dx = \\ = C \cdot S^{\gamma} \left\{ W_3 + h \left[ \left( P_e + \frac{p}{e} \cdot P_p + k \right) r' r \right] \right\},$$

where  $x$  is the density of particles of the shower falling on the absorber,

$W_3 = -3 + 3 \cdot 2^\gamma - 3^\gamma$ ,  $h(z) = -z^\gamma + 3(1+z)^\gamma - 3(2+z)^\gamma + (3+z)^\gamma$ ,  $K \cdot x^{-(\gamma+1)} dx$  is the differential spectrum of densities of registered showers, and  $r' \cdot r \cdot S$  is the effective area of the telescope applied in measurements.

From the ratio  $(TD)/T$  in measurements without absorber we evaluated  $\gamma = 1.41 \pm 0.02$ . By means of formula (2) we can find, for all kinds of coincidences, the quantities  $R(t) = (P_e + p/e \cdot P_p + k)r'r$ , corresponding to different thicknesses of the absorber. From these quantities, after subtracting the penetrating component  $k \cdot r' \cdot r$  and normalisation of the curves obtained for thickness of the absorber  $t=0$ , we evaluate the functions  $R(t) = P_e + p/e \cdot P_p$ . In the normalisation we must take into account the thicknesses of the counter walls.

In this way we obtain from measurements of the group *A* the quantities  $R_1(t)$ ,  $R_2(t)$  and  $R_3(t)$ , corresponding to cases when beneath the absorber there are operated one, two or three trays of counters. The quantity  $R_1(t)$  has been corrected for contribution of penetrating low energy photons. From the values of  $R_1$ ,  $R_2$  and  $R_3$ , we evaluate by means of the formula (1), the ratio  $p/e$  for different thicknesses of the absorber. They are shown in Table I.

We consider a modified anti-coincidence method as being the most convenient.

TABLE I

Values of the ratio  $p/e$  evaluated from measurements

with an absorber placed over the telescope, from the functions				with a modified anti-coincidence method from the differences of rates of coincidences for two positions of the absorber: over the telescope, and between its two upper trays		
$t$ mm.	$R_1(t)$	$R_2(t)$	$R_3(t)$	$(TDEF)$	$(TDE)$	$(TDF)$
3	0.72	0.72	0.66	0.62	0.50	0.61
5	0.82	0.81	0.83	0.98	0.69	0.89
7	0.78	0.78	0.69	0.80	0.77	0.71
10	0.72	0.71	0.68	0.85	0.65	0.86
17	0.61	0.60	0.50	0.42	0.53	0.57
25	0.71	0.72	0.57	0.47	0.91	0.70

In this method we make use of the differences of rates of corresponding coincidences registered by means of an absorber placed over the telescope (group *A*), and between its two upper trays (group *B*). These differences may be expressed by the formula

$$(3) \quad C_{\text{anti}} = A - B = \int_0^{\infty} K \cdot x^{-(\gamma+1)} (1 - e^{-Sx})^3 (1 - e^{-p/e \cdot P_p \cdot r' r S x}) e^{-r S x} dx = \\ = h \left[ r \left( 1 + r' \frac{p}{e} \cdot P_p \right) \right] - h(r).$$

Using this formula we can from experimental data immediately evaluate  $p/e \cdot P_p$ , and, in order to determine the ratio  $p/e$ , we require only one theoretical quantity,  $P_p$ . The main advantage of this method, as compared with the ordinary anti-coincidence method, is that we do not need here any normalisation of the experimental data. The values of the ratio  $p/e$  obtained with the anti-coincidence method are also given in Table I. From this table it is evident that the values of  $p/e$  obtained with both methods are concordant within the limits of error.

In connection with these results let us discuss the following remarks. From the fact that the results of measurements of transition curves obtained in arrangements with different distances of registering trays from the absorber have given concordant values of  $p/e$ , and from the fact that the latter practically do not depend on the thickness of the absorber, we may conclude that our measurements are not influenced markedly by the scattering of particles in the absorber. From the fact of the independence between the results obtained and the thickness of the absorber, we conclude that the result is not greatly influenced, in our case, by the change of mean density of particles of showers registered by our apparatus, which is the result of the change in the thickness of the absorber. In our case the mean density of the particles is comprised in the limits between 10 and 20 per sq. m.

As mentioned above, it is not quite correct to use the values of  $P_e$  and  $P_p$  from Arley's theory. In our case we use only one of these functions, namely  $P_p$ . But this function does not seem to depend strongly on the assumptions of the theory. In this matter the following consideration may help to give a rough picture. From the position of the maximum of the transition curve we can roughly estimate that the main contribution to registering coincidences, besides electrons, comes from photons of the energy of about  $10^3$  eV. The estimation of the average number of electrons beneath the absorber corresponding to one photon over the absorber gives, for  $P_p$ , values between 0.4 and 0.6. Since the statistical error is very small, we assume that the limits given above define roughly the limit of error of the ratio  $p/e$ . So we accept  $p/e = 0.70 \pm 0.15$ , which value is concordant with that given by Milone and concerns the showers of the mean density of particles within the limits of 10 to 20 per sq. m. It is smaller than the value derived from the onedimensional theory of cascades. Most probably it is connected with a different mechanism of scattering of photons and electrons in extensive showers.



The authors wish to express their appreciation of the co-operation of T. Niewodniczański in effecting the measurements and of T. Owsiak for his valuable help in the course of the experiment.

DEPARTMENT OF COSMIC RAYS. NUCLEAR RESEARCH INSTITUTE, POLISH ACADEMY OF SCIENCES, CRACOW

GENERAL PHYSICS LABORATORY, ACADEMY OF MINING AND METALLURGY, CRACOW

## REFERENCES

- [1] C. Millar, *Il Nuovo Cim.* **8** (1951), 279.
- [2] P. Bassi, A. M. Bianchi, C. Manduchi, *Il Nuovo Cim.* **9** (1952), 358.
- [3] C. Milone, *Il Nuovo Cim.* **11** (1954), 241.
- [4] J. M. Massalski, *The transition curve of the photon component of extensive air showers*, *Bull. Acad. Polon. Sci., Cl. III*, **2** (1954), 335.
- [5] N. Arley, *On the theory of the stochastic processes*, 1943.
- [6] K. Greisen, *Phys. Rev.* **75** (1949), 1071.
- [7] L. Jurkiewicz, *On the influence of low energy (2-7 MeV) photons on the absorption curve of extensive air showers in lead*, *Bull. Acad. Polon. Sci., Cl. III*, **2** (1954), 329.



## Hetero-Polyazeotropic Systems. II. Acetonitrile-*n*-Paraffinic Hydrocarbons

by

K. ZIĘBORAK and K. OLSZEWSKI

*Presented by W. ŚWIĘTOSŁAWSKI on September 27, 1956*

### Introductory remarks

In a previous paper [1], the gradual transition from homoazeotropy, through homoazeo-heterozeotropy (azeotropic point lying outside the miscibility gap), to heterozeotropy was investigated in the series of binary systems formed by an azeotropic agent — methanol with the series of *n*-paraffins. In this paper the investigations are described in which the shapes of boiling temperature isobars of binary systems of acetonitrile with *n*-paraffins were examined by using the ebulliometric method. The phenomena observed are different from those observed in the previous investigations. This is due to the fact that the normal boiling temperature of acetonitrile (81.55°C.) is lower than the critical solution temperatures of all the binary systems examined.

Azeotropy in the acetonitrile-hydrocarbon systems has been investigated by Bishop and Denton [2] by the fractional distillation method. They determined the azeotropic parameters, but not the shapes of the boiling temperature isobars and liquid-liquid coexistence curves.

### Experimental part. Preparation of substances

High-grade acetonitrile was distilled from phosphorus pentoxide through a column of 8-10 theoretical plates. Fractions boiling within 0.05° were collected. The hydrocarbons were obtained from synthetic gasoline fractions by treating them manifold with concentrated sulphuric acid and then with soda-solution and water to remove the olefins. The fractions thus obtained were submitted to fractionation in the presence of calcium chloride. Table I contains the refractive indices and



the boiling temperatures of acetonitrile and of very narrow fractions of hydrocarbons, found by means of Świętosławski's ebulliometer.

TABLE I  
Characteristics of the substances

	Boiling temp. °C.	Refractive index $n_D^{20}$
acetonitrile	81.55	1.3447
<i>n</i> -heptane	98.53	1.3875
<i>n</i> -octane	125.7	1.3980
<i>n</i> -nonane	150.0	1.4059
<i>n</i> -decane	174.1	1.4124
<i>n</i> -undecane	195.4	1.4185

The data indicate that the hydrocarbons, especially *n*-nonane and *n*-undecane, contained some amounts of isomers. It seems, however, that their influence on the results was very small.

#### Method of investigation

The boiling and the condensation temperatures were measured in a two-stage semi-micro ebulliometer [3]. Care was taken to avoid any contact of the substances with atmospheric moisture. The atmospheric pressure changes were controlled by the use of a simple ebulliometer filled with water. As the measurements were strictly comparative, the results were corrected to normal pressure. Critical solution temperatures were measured by the conventional Alexieyev method.

#### Results

Boiling temperature isobars, obtained for the binary systems of acetonitrile with hydrocarbons are shown in Fig. 1. The boiling temperature isobar of the acetonitrile-undecane system contains a horizontal section at a temperature higher than the boiling temperature of acetonitrile. The system belongs without any doubt to the heterozeotropic type. The next boiling temperature isobar formed by acetonitrile and *n*-decane has a minimum outside the miscibility curve. The system is therefore homoazeo-heterozeotropic. The isobars formed with *n*-nonane, *n*-octane, and *n*-heptane are characterised by flat minima, typical for heteroazeotropic systems. The data are summarised in Table II.

TABLE II  
Data for the binary systems

System of acetonitrile with:	Boiling temp. of azeotrope or heterozeotrope	Azeotropic depression °C. *)	Type of system	Crit. sol. temp. °C.	Crit. composition in mole per cent of hydrocarbon
<i>n</i> -undecane	81.85	+0.30	heterozeotrope	112.5	23
<i>n</i> -decane	81.45	—0.10	homoazeo-heterozeotrope	107.5	25
<i>n</i> -nonane	79.82	—1.73	heteroazeotrope	100.0	30
<i>n</i> -octane	76.70	—4.85	"	91.5	33
<i>n</i> -heptane	69.55	—12.00	"	84.6	37

\*) The positive sign corresponds to a heterozeotropic system


The results of investigations of mutual solubility of the components are collected in Table II and graphically represented in Fig. 2.

#### Discussion of the results

In the systems examined the transition is observed from heteroazeotropy to heterozeotropy. The mediate stage represents the homoazeo-heterozeotropic system of acetonitrile with *n*-decane. All the systems differ considerably from those formed by methanol with *n*-paraffins [1]. The series fulfils to some extent the assumptions of Stecki who considered the regular solutions showing limited solubility [4]. Probably all the phenomena predicted by the theory would be observed, if additional data for systems with lower boiling hydrocarbons, e. g., *n*-hexane and *n*-pentane or their isomers, were available. The essential difference between the investigated systems and those theoretically considered by Stecki is that the upper critical solution temperatures undergo changes in the real series of systems. In spite of the increase of critical temperatures with increasing molecular weights of the hydrocarbons, the transition phenomena agree with the theory. This is due to the fact that the systems belong to the first class of the classification of heteropolyazeotropic systems, proposed by one of the authors [1].


The data given in Table II indicate that the concentrations in mole per cent of hydrocarbons at the critical temperature decrease linearly with the increasing carbon chain length of the hydrocarbons. The critical temperature increases with the increase of the length of the hydrocarbon molecule. The relative critical temperature increase is the smaller, the larger the hydrocarbon molecule. An exception lies in the difference between the critical solution temperatures of mixtures of acetonitrile

with *n*-heptane and *n*-octane. This is probably due to the presence of some impurities which exert a great influence in that respect.



1. CH<sub>3</sub>CN-C<sub>11</sub>H<sub>24</sub>; 2. CH<sub>3</sub>CN-C<sub>10</sub>H<sub>22</sub>;  
3. CH<sub>3</sub>CN-C<sub>9</sub>H<sub>20</sub>; 4. CH<sub>3</sub>CN-C<sub>8</sub>H<sub>18</sub>;  
5. CH<sub>3</sub>CN-C<sub>7</sub>H<sub>16</sub>;  $p=760$  mm.

Fig. 1. Boiling temperature isobars of binary systems of acetonitrile with *n*-undecane (1), *n*-decane (2), *n*-nonane (3), *n*-octane (4), and *n*-heptane (5). Pressure 760 mm. of Hg



1. CH<sub>3</sub>CN-C<sub>11</sub>H<sub>24</sub>; 2. CH<sub>3</sub>CN-C<sub>10</sub>H<sub>22</sub>;  
3. CH<sub>3</sub>CN-C<sub>9</sub>H<sub>20</sub>; 4. CH<sub>3</sub>CN-C<sub>8</sub>H<sub>18</sub>;  
5. CH<sub>3</sub>CN-C<sub>7</sub>H<sub>16</sub>.

Fig. 2. Mutual solubility curves of the systems of acetonitrile with *n*-undecane (1), *n*-decane (2), *n*-nonane (3), *n*-octane (4), and *n*-heptane (5)



The authors are greatly indebted to Professor W. Świątosławski for his helpful discussion of the phenomena examined.

### Summary

1. The boiling temperature isobars of binary systems formed by acetonitrile with *n*-heptane, *n*-octane, *n*-nonane, *n*-decane and *n*-undecane were determined. Characteristic transition from heteroazeotropy to heterozeotropy was observed.

2. Critical solution temperatures of these systems were measured. Characteristic changes in critical temperatures and compositions have been noticed.

DEPARTMENT OF PHYSICAL CHEMISTRY, WARSAW UNIVERSITY  
RAW MATERIALS DEPARTMENT OF THE INSTITUTE OF PHYSICAL CHEMISTRY,  
POLISH ACADEMY OF SCIENCES

### REFERENCES

- [1] K. Zięborak, Z. Mączyńska, A. Mączyński, *Hetero-polyazeotropic systems I. System methanol-*n*-paraffinic hydrocarbons*, Bull. Acad. Polon. Sci. Cl. III, **4** (1956), 153.
- [2] R. B. Bishop, W. I. Denton, *Ind. Eng. Chem.*, **42** (1950), 883.
- [3] W. Świątosławski, *Ebulliometric Measurements* Reinhold Publ. Corp., New York, 1945.
- [4] J. Stecki, in press.



# TABLE DES MATIÈRES DU VOLUME IV

## Mathématique

J. Albrycht, Some Remarks on the Marcinkiewicz-Orlicz Space . . . . .	1
M. Biernacki, Sur les coefficients tayloriens des fonctions univalentes . . . .	5
W. Pogorzelski, Étude de la solution fondamentale de l'équation parabolique	9
O. Hanš, The Strong Law of Large Numbers for Generalized Random Variables	15
M. Fisz, The Realization of Some Purely Discontinuous Stochastic Processes	63
Z. Szmydt, Sur un nouveau type de problèmes pour un système d'équations différentielles hyperboliques du second ordre à deux variables indépendantes	67
J. Łoś, Abelian Groups that are Direct Summands of Every Abelian Group which Contains Them as a Pure Subgroup . . . . .	73
J. W. Jaworowski and K. Moszyński, A Theorem on Mappings of the Sphere into the Projective Space . . . . .	75
Su-Cheng Chang, On Proper Isomorphism of $(\mu, \Delta, \gamma)$ -Systems. I. . . . .	113
K. Bochenek, Equation $(Vl)^2 = 1$ in the Complex Domain . . . . .	119
A. Plis, On the Estimation of the Existence Domain for Solutions of a Non- Linear Partial Differential Equation of the First Order . . . . .	125
T. Ważewski, Sur la méthode de A. Plis de déterminer le domaine d'existence de la solution du problème de Cauchy pour les équations aux dérivées partielles du premier ordre. . . . .	131
J. Mikusiński, Équations différentielles à coefficients constants considérées dans les espaces linéaires généraux . . . . .	137
J. Łoś, On the Torsion-Free Abelian Groups with Hereditarily Generating Sequences . . . . .	169
L. Włodarski, Propriétés des méthodes continues de limitation du type de Borel	173
L. Włodarski, Sur la concordance entre les méthodes intégrales de sommation du type de Borel . . . . .	177
L. Maurin, Die Lösung im Grossen gemischter Probleme für Systeme inhomogener Linearer Differentialgleichungen . . . . .	179
J. W. Jaworowski, Theorem on Antipodes for Multi-Valued Mappings and a Fixed Point Theorem. . . . .	187
B. Knaster, Sur la fixation des décompositions . . . . .	193
S. Marcus, Sur un problème de la théorie de la mesure de H. Steinhaus et S. Ru- ziewicz . . . . .	197
S. Marcus, Sur un problème F. Hausdorff concernant les fonctions symétriques continues . . . . .	201
A. Schinzel, and Y. Wang, A Note on Some Properties of the Functions $\varphi(n)$ , $\delta(n)$ and $\theta(n)$ . . . . .	207
K. Kuratowski, On a Characterisation of Connected Domains in Locally Con- nected Spaces . . . . .	211
S. Łojasiewicz, Sur la valeur d'une distribution dans un point . . . . .	239
Z. Polniakowski, On Certain Theorems of the Mercer Type . . . . .	243



M. Krzyżański, Sur l'allure asymptotique des solutions de l'équation du type parabolique . . . . .	247
M. Warmus, Calculus of Approximations . . . . .	253
A. Bielecki, Une remarque sur la méthode de Banach-Cacciopoli-Tikhonov dans la théorie des équations différentielles ordinaires . . . . .	261
A. Bielecki, Une remarque sur l'application de la méthode de Banach-Cacciopoli-Tikhonov dans la théorie de l'équation $s=f(x,y,z,p,q)$ . . . . .	265
W. Słowikowski, Note on the Application of the Pauli Ring to form the Metric Tensor in the General Theory of Relativity . . . . .	313
A. Bielecki, Réduction des axiomes de congruence de Hilbert . . . . .	321
P. I. Hilton, Remark on the Tensor Product of Modules . . . . .	325
R. D. Anderson, Some Remarks on Totally Disconnected Sections of Monotone Open Mappings . . . . .	329
K. Borsuk and A. Kosiński, On Connections Between the Homology Properties of a Set and of Its Frontier . . . . .	331
A. Kosiński, On Mappings which Satisfy Certain Conditions on Boundary . . . . .	335
J. Perkal, On the $\varepsilon$ -Length . . . . .	399
A. Hulanicki, Algebraic Characterisation of Abelian Groups which Admits Compact Topologies . . . . .	405
W. Pogorzelski, Propriétés des intégrales de l'équation parabolique générale . . . . .	407
E. Szaśiada, An Application of Kulikov's Basic Subgroups in the Theory of Abelian Mixed Groups . . . . .	411
S. Balcerzyk and Jan Mycielski, On Free Subgroups of Topological Groups . . . . .	415
Jan Mycielski, On the Decompositions of Euclidean Spaces . . . . .	417
Jan Mycielski, On the Congruence of Sets . . . . .	419
J. Mikusiński, Sur la croissance de la fonction opérationnelle $\exp(-s^a\lambda)$ . . . . .	423
S. Mrówka, On the Notion of Completeness in Proximity Spaces . . . . .	477
S. Mrówka, On Universal Spaces . . . . .	479
S. Mrówka, On Quasi-Compact Spaces . . . . .	483
Jan Mycielski, S. Świerczkowski and A. Zięba, On Infinite Positional Games . . . . .	485
R. Sikorski and T. Traczyk, On Some Boolean Algebras . . . . .	489
A. Bielecki, Remarques sur la méthode de T. Ważewski dans l'étude qualitative des équations différentielles ordinaires . . . . .	493
A. Bielecki, Sur une méthode de régularisation des équations différentielles ordinaires dont les intégrales ne remplissent pas la condition d'unicité . . . . .	497
A. Bielecki, Certaines propriétés topologiques des intégrales des équations différentielles ordinaires . . . . .	503
W. Wolibner, Sur le mouvement des corps friables . . . . .	507
A. Lelek, Sur les décompositions en trois continus . . . . .	511
C. Olech, On the Asymptotic Behaviour of the Solutions of a System of Ordinary Non-Linear Differential Equations . . . . .	555
J. Wolska, Problème aux limites à la dérivée tangentielle pour l'équation du type elliptique . . . . .	563
Shao Pin-Tsung, On the Distribution of the Values of a Class of Arithmetical Functions . . . . .	569
Wu Wen-tsün, On the Imbedding of Polyhedrons in Euclidean Spaces . . . . .	573
Z. Szmydt, Sur une généralisation des problèmes classiques concernant un système d'équations différentielles hyperboliques du second ordre à deux variables indépendantes . . . . .	579
S. Świerczkowski, On Cyclic Ordering Relations . . . . .	585
Pan Cheng-Tung, On $\sigma(n)$ and $\varphi(n)$ . . . . .	637
Cheng Min-teh and Cheng Young-hoo, Fractional Integrals of Functions of Several Variables with Applications to the Theory of Approximation . . . . .	639

W. Słowikowski, A Generalisation of Mikusiński's Operational Calculus . . .	645
R. Sikorski, A Theorem on Non-Classical Functional Calculi . . . . .	649
Z. Polniakowski, On Some Tauberian Theorems . . . . .	651
A. T. Bharucha-Reid, On Random Elements in Orlicz Spaces . . . . .	657
R. Engelking, Sur l'impossibilité de définir la limite topologique inférieure à l'aide des opérations dénombrables de l'opération de fermeture . . . . .	659
A. Mostowski, On Models of Axiomatic Set-Theory . . . . .	663
F. Albrecht, Un théorème de comportement asymptotique des solutions des systèmes d'équations différentielles . . . . .	737
A. Pliś, Generalisation of the Cauchy Problem to a System of Partial Differential Equations . . . . .	741
S. Paszkowski, On the Accuracy of Approximation with Nodes . . . . .	745
A. Pliś, Sets Filled by Asymptotic Integrals of Ordinary Differential Equations	749
J. Simaika, Sur une mesure de la dispersion d'une distribution de directions	753
M. Kowalski, On the Determinants of Wroński in Linear Rings . . . . .	789
Yin Wen-Lin, Note on the Representation of Large Integers as Sums of Primes	793
S. C. Chang, On Algebraic Structures and Homotopy Invariants . . . . .	797
H. Steinhaus, Sur la division des Corps matériels en parties . . . . .	801

### Astronomie

K. Rudnicki, Der Einfluss der Bewegung der Dunelmaterie auf den Lichtwechsel der Sterne . . . . .	427
J. Smak, On the Motion of Red Dwarfs with Emission Spectra . . . . .	429
M. Karpowicz, On the Motion of B Stars in the Local System. . . . .	587
M. Karpowicz, On the Rotation of the Local System . . . . .	591

### Physique Théorique

W. Królikowski and J. Rzewuski, On the Equation for a Distinguished Component of the State Vector . . . . .	19
G. Marx, On the Variational Principle of Dielectrics . . . . .	29
G. Marx und K. Nagy, Zum Problem der Impulsübertragung durch un stetige elektromagnetische Wellen in dielektrischen Medien . . . . .	79
W. Czyż and J. Sawicki, Polarization of Nucleons from Photonuclear Reactions	141
W. Kołos, On the Weizsäcker Correction in the Statistical Theory of Molecular Structure . . . . .	147
L. Infeld and I. Plebański, On an Operational Method of Solving the Klein-Gor- don Equation . . . . .	215
W. Kołos, Cross-Section of the Methyl Mercaptan Molecule for the Scattering of Slow Neutrons . . . . .	269
M. Suffczyński, Two-Centre Integrals Over the Atomic Sphere . . . . .	273
K. Nagy, Über die Bewegungsgleichung des magnetischen Dipols . . . . .	341
L. Infeld and J. Plebański, A Simple Derivation of the Equation of Motion in Classical Electrodynamics . . . . .	347
Shoichi Sakata, On a Composite Model for the New Particle . . . . .	353
R. Teisseyre, New Method of Solving the Diffraction Problem for a Dipole Field . . . . .	433
A. Trautman, On a Generalisation of the Einstein—Infeld Approximation Method . . . . .	439
A. Trautman, Solution of One-Body Problem by the Einstein-Infeld Ap- proximation Method . . . . .	443

J. I. Horváth, On the Theory of the Electromagnetic Field in Moving Dielectrics	447
M. Miąsek, Determination of Valence Band in Metallic Sodium by the Parzen Variational Method	453
W. Czyż and J. Sawicki, Polarisation of Nucleons from Photodisintegration of Deuteron. I. Medium Energies	515
P. Zieliński, A Note on Nuclear Isomerism	595
J. Wrzecionko, Character of Interaction Potential Between Electron and Photon	669
A. Trautman, On the Conservation Theorems and Equations of Motion in Covariant Field Theories	675
A. Trautman, Killing's Equations and Conservation Theorem	679
K. Nagy, Über die Bewegungsgleichungen des Pol-Dipol-Teilchens	683
L. Infeld and J. Plebański, On Modified Dirac $\delta$ -Functions	687
L. Infeld and J. Plebański, On a Covariant Formulation of the Equation of Motion	757
L. Infeld and J. Plebański, On the "Dipole Procedure" in General Relativity Theory	763
M. Miąsek, The Calculation of the Matrix Components of Energy for Hexagonal Close-Packed Structure	805

### Physique Expérimentale

Z. Pająk and A. Piekara, Thermal Independence of Permittivity of Ferroelectric Heterogeneous Systems	83
A. Tramer, Raman Spectra of Sulphur Dioxide Solutions	355
Z. Pająk, F. Kaczmarek and J. Jastrzębski, Delay-Effect in Ferroelectric titanates Below the Curie Point	457
B. Moszyńska, Sur l'influence de la température sur la largeur des raies de tetrachlorure de carbone	769
M. Mięslowicz, O. Stanisławski and W. Wolter, Investigation of an Electromagnetic Cascade of Very High Energy in the First Stage of Its Development in Nuclear Emulsions	811
J. Babecki, L. Jurkiewicz, J. M. Massalski and M. Mięslowicz, The Ratio of the Number of Photons to the Number of Electrons in Extensive Air Showers	817

### Chimie

W. Palczewska, The Influence of Polar Organic Substances and Catalytic Poisons on Hydrogen Overvoltage on Nickel and Iron Cathodes	37
M. Śmiałowski, Is There a Quantitative Relationship Between Hydrogen Overvoltage and the Solubility of This Gas in Metals?	43
Z. Kurtyka, The Positive-Negative Azeotrope: Glycol- <i>o</i> -Cresol-2, 4, 6-Col-lidine. XXIII	49
T. Urbański, On Hydrogen Bonds in Some Nitroalcohols	87
W. Świątosławski, K. Zięborak and J. Stecki, Classification of Binary Systems with Limited Mutual Solubility	97
J. Kamecki and Z. Zembura, The Anodic Behaviour of Iron in Phosphoric Acid Solutions	101
J. Dereń, J. Haber and S. Mrowec, Relation Between the Electric Conductivity and Catalytic Activity of ZnO—Cr <sub>2</sub> O <sub>3</sub> Mixed Catalysts	107
T. Urbański and J. Fałęcki, On Nitration of Salicylhydroxamic Acid	151



K. Zięborak, Z. Maczyńska and A. Maczyński, Hetero-Polyazeotropic Systems. I. System Methanol- <i>n</i> -Paraffinic Hydrocarbons . . . . .	153
W. Świętosławski and W. Malesiński, Reduced Equation Characterizing Some Families of Binary Azeotropic Systems . . . . .	159
D. Gürne and T. Urbański, On the Synthesis and Degradation of Certain Derivatives of Tetrahydro-1,3-Oxazine. II . . . . .	221
T. Urbański and B. Szczeciński, On Acetylation of Bis-(2-hydroxymethyl-2-nitro)-butylamine . . . . .	225
K. Gumiński, An Attempt to Elucidate the High Pressures of Hydrogen in Iron Cathodically Saturated with Hydrogen . . . . .	227
A. Kręglewski, The Critical Region of Liquids I. Liquid-Vapour Miscibility and Retrograde Condensation . . . . .	233
J. Michalski and J. Wieczorkowski, The Action of Alkyl Thiocyanates on Esters of Acids of Triavalent Phosphorus Synthesis of O,O,S-Trialkyl thiophosphate . . . . .	279
J. Stecki, Heteroazeotropic Binary Systems. I. Influence of Temperature on the Equilibrium Pressure and on the Vapour Composition . . . . .	283
J. Stecki, Heteroazeotropic Binary Systems. II. Influence of the Pressure and of the Heats of Mixing . . . . .	289
W. Malesiński, Series of Binary Azeotropic Systems Characterised by Asymmetrical Azeotropic Ranges . . . . .	295
W. Malesiński, On the Relation Between the Azeotropic Ranges and the Binary Regular Solution Constants in Three-Component Systems . . . . .	303
T. Urbański and B. Skowrońska-Serafin, Reactions of Aromatic Amines with Cyanguanidine. Reactions of Phenyl Amideurea and Their Derivatives with 2-aminopyridine . . . . .	361
T. Urbański, and B. Skowrońska-Serafin, Reactions of Aromatic Amines with Cyanguanidine. Reactions of Phenylamidineurea and Their Derivatives with <i>N</i> -methylaniline . . . . .	363
W. Malesiński, The Composition of Ternary Positive Azeotropes. I . . . . .	365
W. Malesiński, Boiling Temperatures of Positive Ternary Azeotropes . . . . .	371
T. Urbański, On Hydrogen Bonds in Some Nitroalcohols. II. Infra-Red Absorption Bonds of Nitrogroups . . . . .	381
B. Kamieński and I. Kulawik, The Influence of the Thickness of the Adsorption Layer on the Potential of the Microelement Used in Chromatography . . . . .	387
T. Urbański, Z. Eckstein and R. Wojnowska, On Reaction of 1-Nitromethylcyclohexanol and Its Halogen Derivatives with Formaldehyde and Acetaldehyde . . . . .	461
B. Baranowski, The Seith-Wever Effect Interpretad from the Point of View of Thermodynamics of Irreversible Processes . . . . .	465
J. K. N. Jones, R. Koliński, H. Piotrowska and T. Urbański, On Formation of a Derivative of 1,5 diazabicyclo (3, 3, 3) undekane from 1-nitropropane, Formaldehyde and Ammonia . . . . .	521
W. Malesiński, Composition of Ternary Positive Azeotropes. II. . . . .	523
B. Kamieński and J. Kulawik, The Potential of the Adsorption Micro-electrode in Air Contaminated by Reducing Gases . . . . .	529
A. Bielański, J. Dereń, J. Haber and S. Mrowec, The Influence of the Composition of Zn O-Fe <sub>2</sub> O <sub>3</sub> and of ZnO-Cr <sub>2</sub> O <sub>3</sub> Mixed Catalysts on Their Electric Conductivity and Catalytic Activity . . . . .	533
Z. Ruziewicz, Luminescence of Oxide Films Produced During Anodic Oxidation of Aluminium . . . . .	537
Z. Ruziewicz, Luminescence of Aluminium Electrodes During Electrolysis by Alternating Current in Solutions of Oxalic Acid . . . . .	543

T. Urbański, On a New Colour Reaction of Aldehydes . . . . .	599
J. Niemiec and W. Trzebiatowski, The Existence of $\sigma$ - and $\chi$ - Phases in Rhenium Alloys . . . . .	601
A. Gołębiewski, Conductance of Mono- and Poly-Crystalline Resorcinol . . .	605
W. Malesiński, Boiling Temperature of Positive Ternary Azeotropes. II . . .	609
T. Penkala, Influence of Ion Size on Quantitative Proportions of the Components in Binary Salts. IV . . . . .	615
T. Penkala, Gradual Transition in Series of Binary Systems from Compounds Melting Incongruently to Congruently. V . . . . .	619
W. Trąbczyński, Binary Azeotropes Formed by Propionic Acid with the Series of <i>n</i> -Paraffins . . . . .	623
Z. Eckstein and T. Urbański, On Fungicide Activities of Some Aryloxyacetyldroxyamic Acids . . . . .	627
W. Świątosławski and W. Malesiński, Azeotropic Depression as a Function of the Composition of Series of Ternary Positive Azeotropes . . . . .	693
W. Malesiński, Composition of Positive, Saddle and Negative Ternary Homoazeotropes . . . . .	701
W. Malesiński, Boiling Temperatures of Positive Saddle, and Negative Ternary Homoazeotropes . . . . .	709
J. Sędzimir and M. Pawełkova, Kinetics and Mechanism of Cementation Processes. I. Model Lead-Zinc Cementation Cell . . . . .	717
J. Sędzimir, Kinetics and Mechanism of Cementation Processes. II. Effect of Cementation Products on Kinetics of the Process . . . . .	723
T. Urbański and K. Szye-Lewańska, Some Remarks on the Structure of Aniline Black on the Basis of Ultraviolet Absorption. Spectra . . . . .	775
J. H. Kolutowska, On the Production of Sodium Pyrophosphite $\text{Na}_2\text{H}_2\text{P}_2\text{O}_6$ by Pyrolysis of Disodium Hypophosphate $\text{N}_2\text{H}_2\text{P}_2\text{O}_6$ . . . . .	783
K. Zięborak and K. Olszewski, Hetero-Polyazeotropic Systems. II. Acetonitrile- <i>n</i> -Paraffinic Hydrocarbons . . . . .	823

### Géologie

S. Z. Różycki, New Stratigraphic Division of the Dogger, Based on Ammonite Fauna, in the Marginal Areas of the Holy Cross Mountains . . . . .	55
H. Teisseyre, Some Remarks on the Tectonic Structure of Caledonides and Variscides in the Sudeten . . . . .	165
L. Koszarski, Observations on the Sedimentation of the Cieżkowice Sandstone near Cieżkowice (Carpathian Flysch) . . . . .	393
R. Unrug, Preferred Orientation of Pebbles in Recent Gravels of the Dunajec River in the Western Carpathians . . . . .	469
B. Tomeczyk and L. Teller, The Ludlow Deposits in Eastern Poland . . . .	549
S. Bukowy, Observations on the Sedimentation of the Babica Clays (Paleocene) in the Meddle Carpathians . . . . .	631
K. Birkenmajer, Sedimentary Characteristics of the Jarmuta Beds (Maestrichian) in the Pieniny Klippen Belt (Central Carpathians) . . . . .	729

# LISTE DES AUTEURS DU VOLUME IV

Albrecht F. . . . .	741	Kołos W. . . . .	147, 269
Albrycht J. . . . .	1	Kosiński A. . . . .	331, 335
Anderson R. D. . . . .	329	Koszarski L. . . . .	393
Babecki J. . . . .	817	Kowalski M. . . . .	789
Balcerzyk S. . . . .	415	Kręglewski A. . . . .	233
Baranowski B. . . . .	465	Królikowski W. . . . .	19
Bielański A. . . . .	533	Krzyżański M. . . . .	247
Bielecki A. 261, 265, 321, 493, 497, 503		Kulawik I. . . . .	387, 529
Biernacki M. . . . .	5	Kuratowski K. . . . .	211
Birkenmajer K. . . . .	731	Kurtyka Z. . . . .	49
Bharucha-Reid A. T. . . . .	657	Lelek A. . . . .	511
Bochenek K. . . . .	119	Łojasiewicz S. . . . .	239
Borsuk K. . . . .	331	Łoś J. . . . .	73, 169
Bukowy S. . . . .	631	Marcus S. . . . .	197, 201
Chang S. C. . . . .	113, 797	Marx G. . . . .	29, 79
Chen Yung-hoo . . . . .	641	Malesiński W. . 159, 295, 303, 365, 371	
Cheng Min-teh . . . . .	641		523, 609, 695, 703, 711
Czyż W. . . . .	141, 515	Maurin L. . . . .	179
Dereń J. . . . .	107, 533	Mączyńska Z. . . . .	153
Eckstein Z. . . . .	461, 627	Mączyński A. . . . .	153
Engelking R. . . . .	661	Massalski J. M. . . . .	817
Fałęcki J. . . . .	151	Miąsek M. . . . .	453, 805
Fisz M. . . . .	63	Michalski J. . . . .	279
Gołębiewski A. . . . .	605	Mięsowicz M. . . . .	811, 817
Gumiński K. . . . .	227	Mikusiński J. . . . .	137, 423
Gürne D. . . . .	221	Mostowski A. . . . .	665
Haber J. . . . .	107, 533	Moszyńska B. . . . .	773
Hanš O. . . . .	15	Moszyński K. . . . .	75
Hilton P. J. . . . .	325	Mrowec S. . . . .	107, 533
Horváth J. I. . . . .	447	Mrówka S. . . . .	477, 479, 483
Hulanicki A. . . . .	405	Mycielski J. . . . .	415, 417, 419, 485
Infeld L. . . . .	215, 347, 689, 761, 767	Nagy K. . . . .	79, 341, 685
Jastrzębski J. . . . .	457	Niemiec J. . . . .	601
Jaworowski J. W. . . . .	75, 187	Olech C. . . . .	555
Jones J. K. N. . . . .	521	Olszewski K. . . . .	827
Jurkiewicz L. . . . .	817	Pająk Z. . . . .	83, 457
Kaczmarek F. . . . .	457	Palczewska W. . . . .	37
Kamecki J. . . . .	101	Pan Cheng-Tung . . . . .	639
Kamiński B. . . . .	387, 529	Paszkowski S. . . . .	749
Karpowicz M. . . . .	587, 591	Pawełkowa M. . . . .	719
Knaster B. . . . .	193	Penkala T. . . . .	615, 619
Koliński R. . . . .	521	Perkal J. . . . .	399
Kolitowska J. H. . . . .	787	Piekara A. . . . .	83

Piotrowska H. . . . .	521	Świętosławski W. . . . .	97, 159, 695
Plebański J. . . . .	215, 347, 689, 761, 767	Teisseyre H. . . . .	165
Pliś A. . . . .	125, 745, 753	Teisseyre R. . . . .	433
Pogorzelski W. . . . .	9, 407	Teller L. . . . .	549
Polniakowski Z. . . . .	243, 653	Tomeczyk H. . . . .	549
Różycki S. Z. . . . .	55	Traczyk T. . . . .	489
Rudnicki K. . . . .	427	Tramer A. . . . .	355
Ruziewicz Z. . . . .	537, 543	Trautman A. . . . .	439, 443, 677, 681
Rzewuski J. . . . .	19	Trąbceżyński W. . . . .	623
Sakata S. . . . .	353	Trzebiatowski W. . . . .	601
Sawicki J. . . . .	141, 515	Unrug R. . . . .	469
Sasiada E. . . . .	411	Urbański T. 87, 151, 221, 225, 361, 363, 381, 461, 521, 599, 627, 779	
Schinzel A. . . . .	207	Wang Y. . . . .	207
Sędzimir J. . . . .	719, 725	Warmus M. . . . .	253
Shao Pin-Tsung. . . . .	569	Ważewski T. . . . .	131
Sikorski R. . . . .	489, 651	Wieczorkowski J. . . . .	279
Simaika J. . . . .	757	Włodarski L. . . . .	173, 177
Skowrońska-Serafin B. . . . .	361, 363	Wojnowska H. . . . .	461
Słowikowski W. . . . .	313, 645	Wolibner W. . . . .	507
Smak J. . . . .	429	Wolska J. . . . .	563
Stanisz O. . . . .	811	Wolter W. . . . .	811
Stecki J. . . . .	97, 283, 289	Wrzecionko J. . . . .	671
Steinhaus H. . . . .	801	Wu Wen-tsün . . . . .	573
Suffczyński M. . . . .	273	Yin Wen-Lin . . . . .	793
Szczyński B. . . . .	225	Zembura Z. . . . .	101
Szmydt Z. . . . .	67, 579	Zieliński P. . . . .	595
Szyc-Lewańska K. . . . .	779	Zięba A. . . . .	485
Śmiałowski M. . . . .	43	Zięborak K. . . . .	97, 153, 823
Świerczkowski S. . . . .	485, 585		



# ERRATA AU VOLUME IV

Page/ligne	Au lieu de	Lire
1 <sup>6</sup>	$M$	$\overline{M}$
3 <sup>7</sup>	$ u p/P, \quad  v q/Q$	$ u p/p, \quad  v q/q$
110	Le texte indiqué sous la Fig. 3 se rapporte à la Fig. 4, et vice versa.	
207 <sub>13</sub>	$A_1$	$A_i$
207 <sub>8</sub>	<i>positive</i>	<i>integral</i>
208 <sup>1</sup>	positive number	integer
208 <sup>6</sup>	$M_Z(x)$	$M_Z(X)$
240 <sup>3</sup>	$+0( x-x_0 )^n,$	$+0( a-x_0 ^n),$
241 <sub>4</sub>	$1/ x ^a \sin 1/ x ^\beta$	$ x ^{-a} \sin  x ^{-\beta}$
241 <sub>4</sub>	intégrales	intégrables
245 <sup>11</sup>	$\sum_{v=0}^k$	$\sum_{v=0}^l$
251 <sup>8,9,15</sup>	Rayer $t$ dans les équations.	
251 <sup>18</sup>	Rayer $t$ du côté gauche de l'équation.	
314 <sup>11</sup>	$a \in \mathfrak{P}$	$\mathbf{a} \in \mathfrak{P}$
316 <sup>2</sup>	$\mathbf{P}$	$\mathfrak{P}$
317 <sup>10</sup>	$\Gamma_{ki}^i \mathbf{df}(\delta_k \mathbf{g}_i, \mathbf{g}^i) \left( \frac{\delta \mathbf{g}_i}{\delta \mathbf{x}^k}, g^i \right)$	$\Gamma_{ki}^i \mathbf{df}(\delta_k \mathbf{g}_i, \mathbf{g}^i)$
317 <sup>12-13</sup>	$\sqrt{R_{kln}^i} = 0,$	$R_{kln}^i = 0,$
318 <sup>12</sup>	$D\mathbf{x}^* = \mathbf{g}^{i_0 i_1 \dots i_n} \dots$	$D\mathbf{x}^* = (n+1) \mathbf{g}^{i_0 i_1 \dots i_n} \dots$
318 <sup>13</sup>	$= \mathbf{g}_{i_0 i_1 \dots i_n}$	$= (n+1) \mathbf{g}_{i_0 i_1 \dots i_n}$
319 <sup>22</sup>	$D^* \mathbf{a} = \mu f$	$D^* \mathbf{a} = \mu \mathbf{f}^*$
319 <sup>21-22</sup>	$a = a^*$	$\mathbf{a} = \overline{\mathbf{a}}$
322 <sub>13</sub>	$\triangle ABC \cong \triangle ABC',$	$\triangle ADC \cong \triangle ADC'$
322 <sub>7</sub>	$\triangle ACB \cong \triangle AOB,$	$\triangle ACD \cong \triangle ACD$
323 <sup>7</sup>	$AB$	$AC$
323 <sup>20</sup>	l'angle droit	l'angle droit $A$
405 <sup>6</sup>	gives a $\varphi$ algebraic	gives an algebraic
405 <sub>5</sub>	$G = D + \sum_u^* J_n / \sum_j^* J_j,$	$G = D + \sum_a^* J_{pa} / \sum_\beta^* J_{p\beta}$
428 <sup>1</sup>	$\cos^2 \lambda$	$\sin^2 \lambda$

Page/ligne	Au lieu de	Lire
428 <sup>15</sup>	$\cos^2 \lambda$	$\sin^2 \lambda$
428 <sup>21</sup>	$7 \cdot 10^{-14}$	$6 \cdot 10^{-18}$
442 <sub>7</sub>	$-\frac{1}{2} g_0^2 g_{00;ik}$	$-\frac{1}{2} g_{00;ik}$
454 <sup>17</sup>	$8 O_{1s,2}^2$	$8 O_{1s,1}^2$
455 <sup>10</sup>	$I_0=0$	$\delta I_0=0$
494 <sup>23</sup>	$O(R)$	$O(Q)$
500 <sup>3</sup>	$0 < \eta(x) \leq \delta(x)$	$0 < \eta(t,x) \leq \delta(t,x)$



